

## Repellent activity of *Glycosmis* plant extracts against two stored product insects

[Actividad repelente de extractos de plantas *Glycosmis* contra dos insectos de productos almacenados]

Kai YANG<sup>1</sup>, Shan Shan GUO<sup>1</sup>, Wen Juan ZHANG<sup>1</sup>, Cheng Fang WANG<sup>1,2</sup>, Jiao HAN<sup>3</sup>, Zhu Feng GENG<sup>3</sup>, Yan WU<sup>1,4</sup>, Shu Shan DU<sup>1</sup> & Zhi Wei DENG<sup>3</sup>

<sup>1</sup>Beijing Key Laboratory of Traditional Chinese Medicine Protection and Utilization, Beijing Normal University, No. 19, Xijiekouwai Street, Beijing 100875, China

<sup>2</sup>China CDC Key Laboratory of Radiological Protection and Nuclear Emergency, National Institute for Radiological Protection, Chinese Center for Disease Control and Prevention, Xicheng District, Beijing 100088, China

<sup>3</sup>Analytical and Testing Center, Beijing Normal University, Beijing 100875, China

<sup>4</sup>Technical Center of China Tobacco Guangxi Industrial Co., Ltd., Nanning 530001, China

Contactos / Contacts: Shu Shan DU - E-mail address: [dushushan@bnu.edu.cn](mailto:dushushan@bnu.edu.cn)

Contactos / Contacts: Cheng Fang WANG - E-mail address: [wangchengfang@mail.bnu.edu.cn](mailto:wangchengfang@mail.bnu.edu.cn)

**Abstract:** In the present study, the repellent activities of the leaf and/or stem crude extracts of *Glycosmis lucida* Wall. ex Huang, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha* Huang, *G. pentaphylla* (Retz) Correa. and *G. esquirolii* (Levl.) Tanaka were analyzed by using assays on petri dishes against *Tribolium castaneum* and *Liposcelis bostrychophila*. The leaf and stem extracts of *G. lucida*, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha* and *G. esquirolii* possessed significant repellent activities against *T. castaneum*, the same level repellent with the positive control, DEET. However, the extracts of *G. pentaphylla*, no repellency but some insect attractant was observed. Moreover, they also showed repellent activities against *L. bostrychophila*. These results indicate that extracts from *G. lucida* and *G. oligantha* leaf could be a source of novel repellent against insects.

**Keywords:** extraction, *Glycosmis*, repellency, *Tribolium castaneum*, *Liposcelis bostrychophila*

**Resumen:** En el presente estudio, las actividades repelentes de la hoja y/o tronco de los extractos crudos de *Glycosmis lucida* Wall. ex Huang, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha* Huang, *G. pentaphylla* (Retz) Correa y *G. esquirolii* (Levl.) Tanaka se analizaron mediante el uso de ensayos en placas de Petri contra *Tribolium castaneum* y *Liposcelis bostrychophila*. Los extractos de las hojas y tallo de *G. lucida*, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha* y *G. esquirolii* poseían actividades repelentes significativas contra *T. castaneum*, el mismo nivel repelente del control positivo, el DEET. Sin embargo, los extractos de *G. pentaphylla*, no se observó la repelencia pero sí actividad atrayente de insectos. Por otra parte, también se mostraron las actividades repelentes contra *L. bostrychophila*. Estos resultados indican que los extractos de hojas de *G. lucida* y *G. oligantha* podrían ser una fuente de repelente contra los insectos.

**Palabras clave:** Extracción, *Glycosmis*, repelencia, *Tribolium castaneum*, *Liposcelis bostrychophila*.

Recibido | Received: February 13, 2015

Aceptado | Accepted: May 19, 2015

Aceptado en versión corregida | Accepted in revised form: September 22, 2015

Publicado en línea | Published online: November 30, 2015

**Declaración de intereses | Declaration of interests:** This project was supported by State Key Laboratory of Earth Surface Processes and Resource Ecology, National Natural Science Foundation of China (N° 81374069) and Beijing Municipal Natural Science Foundation (N° 7142093).

**Este artículo puede ser citado como / This article must be cited as:** K Yang, SS Guo, WJ Zhang, CF Wang, J Han, ZF Geng, Y Wu, SS Du, ZW Deng. 2015. Repellent activity of *Glycosmis* plant extracts against two stored product insects. *Bol Latinoam Caribe Plant Med Aromat* 14 (6): 462 – 469.

## INTRODUCTION

Antagonistic storage has been used as one of traditional Chinese medicinal materials conservation methods. It mainly utilizes some traditional Chinese medicinal materials having special volatile odor to store with medicinal materials vulnerable to insects, so as to prevent the insects. According to our survey of literatures, many cases which use this method to prevent the insects were reported (Xia *et al.*, 2000; Wang, 2003; Han *et al.*, 2004; Luo *et al.*, 2012). In order to inherit and develop the traditional method of prevention and control of stored grain insects, we took five *Glycosmis* species as research object and two traditional Chinese medicinal materials storage insects (*Tribolium castaneum* and *Liposcelis bostrychophila*) as the target insects. It was expected that this research work would provide some theoretical basis for the conception of antagonistic storage.

The red flour beetle (*Tribolium castaneum* Herbst) (Coleoptera: Tenebrionidae) is world-widely distributed and among the most economically important stored product pests (Garcia *et al.*, 2005). Infestations can also result in elevated temperature and moisture conditions that lead to an accelerated growth of molds, including toxigenic species (Magan *et al.*, 2003). The booklouse (*Liposcelis bostrychophila* Badonnel) (Psocoptera) is an important emerging pest of stored products. It feed on a wide variety of foods, but they are particularly associated with amylaceous products, such as grain and flour (Nayak, 2006; Throne *et al.*, 2006; Zhou *et al.*, 2012). It is widely distributed in several tropical and sub-tropical countries in Asia, Europe, North America, South America, Africa and Australia (Turner, 1999; Li *et al.*, 2011). This insect's parthenogenic mode of reproduction coupled with its short life cycle in favourable conditions make it particularly troublesome, as it can rapidly infest susceptible commodities (Fisher, 1985; Mills *et al.*, 1992; Phillips & Throne, 2010). Infestations of stored product insects could be controlled by insecticidal treatment of commodities and surfaces (Zettler & Arthur, 2000). However, increasing attention is being given to the development of more ecologically and economically sustainable control methods for stored-product pests, mainly because of the problems already mentioned, for example, the management of the habitat and the use of semiochemical-baited traps, repellents or of biological control (Phillips & Throne, 2010).

It is well known that Rutaceae are used in many countries for their repellent properties against several insects (Venkatachalam & Jebanesan, 2001; Ngassoum *et al.*, 2007; Fogang *et al.*, 2012; Mehmood *et al.*, 2012; Conti *et al.*, 2013; Ali *et al.*, 2013). The *Glycosmis* genus is a species of shrub or dungarunga of the Rutaceae family and distributes in south and southeast of Aisa, northeast of Australia and mainly located in south of nanling, south of Yunnan and southeast of Tibet in China (Huang, 1997). A survey of literature has shown that the chemical composition of genus *Glycosmis* possessed the insecticide activities against several insects (Greger *et al.*, 1996; Latha & Joseph, 1999). Among the *Glycosmis* genus, the repellency of *G. pentaphylla* essential oil against the insect was evaluated (Khan *et al.*, 1983; Pandey *et al.*, 2013). However, little is known about the effectiveness of other *Glycosmis* species extract against stored product insects.

To our knowledge, nothing was reported on the repellent activity of *G. lucida*, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha*, *G. pentaphylla* and *G. esquirolii* extracts against *T. castaneum* and *L. bostrychophila*. In this study, their repellent activity against the red flour beetle and the booklouse were investigated for the first time.

## MATERIAL AND METHODS

### Plant materials

The leaf and stem samples of *G. lucida*, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha*, *G. pentaphylla* and *G. esquirolii* plants were collected in Xishuangbanna, Yunnan province, China (northern latitude: 21°08'~22°36'; east longitude: 99°56'~101°50'), between 2011 June and 2013 June. The specimens were identified by Dr. LX Zhang, Yunnan branch of the Institute of Medicinal Plant Development (IMPLAD), Chinese Academy of Medical Sciences, Yunnan, China, and the voucher specimens were deposited at the Herbarium (BNU) of the College of Resources Science and Technology, Beijing Normal University.

### Extraction

The plant samples were dried naturally for a week. The dried plant samples (100 g) were ground to powders using a grinding mill (High speed grinder 6202, Beijing huanya tianyuan machinery technology co. LTD, Beijing, China) and extracted thrice with methanol (800 mL, 30 min each) at room temperature by using ultrasonic irradiation. All extracts were filtered through Whatman N° 1 filter paper,

evaporated to dryness through rotary evaporator at 45° C under vacuum and further dried under high vacuum overnight. The dried extracts were stored in parafilm glass vials at 4° C until required for analysis.

### Insects

The red flour beetle, *T. castaneum* was obtained from laboratory cultures maintained for the last 2 years in the dark, in incubators at 28-30° and 70-80% relative humidity. The beetles were reared on wheat flour mixed with yeast (10:1, w/w) at 12-13% moisture content. Unsexed adult beetles used in the experiments were about 2 weeks old.

The booklouse, *L. bostrychophila* was obtained from laboratory cultures maintained for the last 2 years in the dark, in incubators at 29-30° and 70-80% relative humidity and was reared on a 1:1:1 mixture, by mass, of milk powder, active yeast and flour. All containers housing insects and the Petri dishes used in experiments were made escape proof with a coating of polytetrafluoroethylene (Fluon, Sino-Rich, Beijing, China). Insects used in the experiments were 1 week old.

### Repellent activity bioassay

The repellent effects of the essential oil against *T. castaneum* and *L. bostrychophila* were assessed by

using assays on petri dishes (Chaubey, 2007). Petri dishes 9 cm in diameter were used to confine beetles during the experiment. The extracts of *G. lucida*, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha*, *G. pentaphylla* and *G. esquirolii* were prepared in ethanol (157.26, 31.45, 6.29 nL/cm<sup>2</sup>), and absolute ethanol was used as the control. Filter paper 9 cm in diameter was cut in half and 500 µL of each concentration was applied separately to half of the filter paper as uniformly as possible with a micropipette. The other half (control) was treated with 500 µL of absolute ethanol. Both the treated half and the control half were then air-dried to evaporate the solvent completely (2 min). A full disk was carefully remade by attaching the tested half to the negative control half with tape. Each reassembled filter paper after treatment with solid glue was placed in a petri dish with the seam oriented in one of four randomly selected different directions to avoid any insecticidal stimuli affecting the distribution of insects. Twenty insects were released in the center of each filter paper disk, and a cover was placed over the petri dish. Five replicates were used, and the experiment was repeated three times. Counts of the insects present on each strip were made after 2 and 4 h. The percent repellency (PR) of each volatile oil was then calculated using the formula:

$$PR (\%) = [(N_c - N_t) / (N_c + N_t)] \times 100$$

Where  $N_c$  is the number of insects present in the negative control half and  $N_t$  is the number of insects present in the treated half. The averages were then assigned to different classes (0 to V) using the following scale (percentage repellency) (Liu & Ho, 1999). Class, % repellency: 0, >0.01 to <0.1; I, 0.1-20.0; II, 20.1-40.0; III, 40.1-60.0; IV, 60.1-80.0; and V, 80.1-100.

As for the booklouse, petri dishes and filter papers were changed to 6 cm in diameter and the concentration of the oil used in the experiments were 126.32, 25.26, 5.05 nL/cm<sup>2</sup>. The half filter paper was treated with 150 µL of the solution. As a positive control, a commercial repellent DEET (*N*, *N*-diethyl-3-methylbenzamide), was used under the conditions as the oil. Analysis of variance (ANOVA) and LSD test were conducted by using SPSS statistics 20 for Windows 2007. Percentage was subjected to an arcsine square-root transformation before ANOVA

and LSD tests.

### RESULTS AND DISCUSSION

Methanol extracts were obtained before carrying out each specific biological assay. The plant extracts yields of six *Glycosmis* species were 1.84% to 7.80% (Table 1). The repellent activity of the leaf and/or stem extracts of medicinal plants against *T. castaneum* and *L. bostrychophila* was determined by using assays on petri dishes (Table 2 and Table 3).

According to our results, except for *G. pentaphylla*, other five *Glycosmis* species extracts possessed strong repellency against *T. castaneum*, especially, the *G. lucida* leaf extract displayed the highest repellency against *T. castaneum* (Table 2). When at the lowest tested concentrations of 6.29 nL/cm<sup>2</sup>, the extract of *G. lucida* leaf still showed stronger repellency (94% and 84%, class V) than the positive control, DEET (78% and 68%, class IV) at 2

h and 4 h after exposure (Table 2). The extracts of *G. oligantha* leaf, *G. esquirolii* leaf displayed the same level repellency against *T. castaneum* compared with the positive control. When at the concentration of 31.45 nL/cm<sup>2</sup>, the extracts of *G. craibii* var. *glabra* leaf showed strong repellency (88%, class V) at 2 h after exposure and the *G. craibii* Tanaka leaf extracts also exhibited strong repellency (90% and 88%, class

V) against *T. castaneum* at 2 h and 4 h after exposure (Table 2). With the concentration decreasing, the repellency of the *Glycosmis* species extracts also weakened gradually. However, the extracts of *G. pentaphylla* leaf and stem especially the *G. pentaphylla* stem, no repellency but some insect attractant properties against the red flour beetle were observed at 2 and 4 h after exposure (Table 2).

**Table 1**  
Methanol extracts of the 6 *Glycosmis* species

Species	Source	Yield (%)	Family
<i>G. lucida</i>	Leaf	5.48	Rutaceae
<i>G. craibii</i> var. <i>glabra</i>	Leaf	7.80	Rutaceae
<i>G. craibii</i> Tanaka	Leaf	1.84	Rutaceae
<i>G. oligantha</i>	Leaf	2.66	Rutaceae
	Stem	2.86	
<i>G. pentaphylla</i>	Leaf	2.79	Rutaceae
	Stem	4.34	
<i>G. esquirolii</i>	Leaf	6.22	Rutaceae
	Stem	5.08	

At the same time, the extracts of the six *Glycosmis* species also possessed repellency against *L. bostrychophila*. When at the highest concentration of 126.32 nL/cm<sup>2</sup>, the extracts of *G. lucida* leaf, *G. oligantha* leaf possessed the same level repellency against *L. bostrychophila* at 2 h after exposure and the extracts of *G. craibii* Tanaka leaf, *G. oligantha* leaf, *G. pentaphylla* leaf, *G. esquirolii* leaf also displayed strong repellency against *L. bostrychophila* at 4 h after exposure (Table 3). While other *Glycosmis* species extracts possessed moderate repellency at 2 h and 4 h after exposure. With the concentration decreasing, the repellency of the six *Glycosmis* species extracts weakened gradually. Compared with the positive control, at the concentrations of 25.26, 5.05 nL/cm<sup>2</sup>, the extracts of all species displayed lower level repellency against *L.*

*bostrychophila*. At the concentration of 5.05 nL/cm<sup>2</sup>, the extracts of *G. pentaphylla* leaf and *G. esquirolii* stem, also no repellency but some insect attractant properties against the booklouse were observed at 4 h after exposure (Table 3). Some extracts were evaluated for repellency against insects (Lou et al., 2013). In China, the extracts derived from Chinese medicinal herbs were also evaluated for insecticidal activity and repellency against the red flour beetle and other insects (e.g., Huang et al., 2007; Yao et al., 2007; Tang et al., 2009a; Tang et al., 2009b; Tang et al., 2009c; Huang et al., 2010; Wei et al., 2014). However, no reports are on bioactivity to the booklouse. In this article, the extracts of 6 *Glycosmis* species were evaluated for repellency against the two grain storage insects, *T. castaneum* and *L. bostrychophila*.

Table 2  
Percentage repellency of the *Glycosmis* species extracts against  
*Tribolium castaneum* after two exposure times

Treatment	Plant part	2h			4h		
		157.26*	31.45*	6.29*	157.26*	31.45*	6.29*
<i>G. lucida</i>	leaf	90±6ab V	86±10abc V	94±6ab V	88±9ab V	76±21ab IV	84±13a V
<i>G. craibii</i> var. <i>glabra</i>	leaf	94±9ab V	88±10ab V	60±19cd III	100±0a V	74±16ab IV	58±14bc III
<i>G. craibii</i> Tanaka	leaf	58±18c III	90±8ab V	60±18cd III	72±20c IV	88±11a V	34±19c II
<i>G. oligantha</i>	leaf	80±15bc IV	76±16bc IV	76±21bc IV	86±14b V	68±18b IV	64±23b IV
	stem	84±14ab V	82±18abc V	42±21d III	92±9ab V	72±16ab IV	30±21c II
<i>G. pentaphylla</i>	leaf	54±19c III	76±16bc IV	-58±17e -	48±21d III	66±19b IV	-8±13d -
	stem	-36±20d -	-62±16d -	-76±15e -	-64±12e -	-76±21c -	-84±10e -
<i>G. esquirolii</i>	leaf	82±18ab V	68±15c IV	64±9cd IV	76±15c IV	90±12a V	42±11bc III
	stem	78±18bc IV	88±18ab V	52±12d III	80±13bc IV	80±8a IV	44±17bc III
DEET		100±0a V	98±3a V	78±14bc IV	96±3a V	82±8a V	68±5b IV

\* Concentration (nL/cm<sup>2</sup>);

a-e Means in the same column followed by the same letters do not differ significantly ( $P < 0.05$ ) in ANOVA and LSD tests. PR was subjected to an arcsine square-root transformation before ANOVA and LSD tests.

To our knowledge, the repellent activity of *G. lucida*, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha*, *G. pentaphylla* and *G. esquirolii* has not been reported before. Previous studies have isolated carbazole alkaloid, acridone alkaloids from *G. pentaphylla* (Govindachari et al., 1966; Yang et al., 2012), hydroquinone diglycoside acyl esters, isoflavone diglycosides from *G. pentaphylla* (Wang et al., 2006a; Wang et al., 2006b) and flavanones, flavanonols from *Glycosmis* species (Lukaseder et al., 2009). Some compounds from *Glycosmis* species with antitumor, antifungal, antibacterial and insecticidal have been observed (Bhattacharyya et al., 1985; Greger et al., 1996; Quader et al., 1999; Ito et al., 2004; Sripisut et al., 2012). However, their association with repellency activity is currently

unknown. The results pointed out the potential value of the compounds from the extracts used to prevent the stored products insects.

## CONCLUSIONS

The results have shown that the extracts of *G. lucida*, *G. craibii* var. *glabra*, *G. craibii* Tanaka, *G. oligantha* and *G. esquirolii* exhibit strong repellent activity against *T. castaneum*. They show potential to be developed as possible natural repellent for control of stored product insects. However, for the practical application of the extracts, further studies on the safety of the extracts to humans and on the effect of the compounds from the plant extracts are necessary to improve the efficacy.

Table 3  
Percentage repellency of the *Glycosmis* species extracts against *Liposcelis bostrychophila* after two exposure times

Treatment	Plant part	2h			4h		
		126.32*	25.26*	5.05*	126.32*	25.26*	5.05*
<i>G. lucida</i>	leaf	84±10ab	58±8b	34±12bc	76±6cd	48±18bc	20±12cd
		V	III	II	IV	III	I
<i>G. craibii</i> var. <i>glabra</i>	leaf	78±17bc	70±8ab	38±11bc	78±10bcd	68±12abc	18±13cd
		IV	IV	II	IV	IV	I
<i>G. craibii</i> Tanaka	leaf	76±14bc	66±13ab	34±11bc	86±13bc	76±13ab	46±12bc
		IV	IV	II	V	IV	III
<i>G. oligantha</i>	leaf	88±8ab	58±11b	40±12bc	100±5a	74±11ab	32±18c
		V	III	II	V	IV	II
	stem	80±9abc	62±15ab	48±8bc	62±12d	56±15bc	60±8b
<i>G. pentaphylla</i>	leaf	70±17bc	52±6b	16±19de	84±14bc	58±15bc	-24±24e
		IV	III	I	V	III	-
	stem	70±13bc	50±7b	50±9bc	76±10cd	38±6c	10±14cd
<i>G. esquirolii</i>	leaf	74±12bc	54±13b	68±17ab	88±10bc	76±13ab	36±18bc
		IV	III	IV	V	IV	II
	stem	66±13c	28±20c	6±7e	58±14d	38±8c	-6±12de
DEET		94±6a	82±5a	86±8a	92±5ab	84±3a	82±8a
		V	V	V	V	V	V

\* Concentration (nL/cm<sup>2</sup>);

a-e Means in the same column followed by the same letters do not differ significantly ( $P < 0.05$ ) in ANOVA and LSD tests. PR was subjected to an arcsine square-root transformation before ANOVA and LSD tests.

#### ACKNOWLEDGEMENTS

This project was supported by State Key Laboratory of Earth Surface Processes and Resource Ecology, National Natural Science Foundation of China (N° 81374069) and Beijing Municipal Natural Science Foundation (N° 7142093).

#### REFERENCES

- Ali A, Demirci B, Kiyani HT, Bernier UR, Tsikolia M, Wedge DE, Khan IA, Baser KHC, Tabanca N. 2013. Biting deterrence, repellency, and larvicidal activity of *Ruta chalepensis* (Sapindales: Rutaceae) essential oil and its major individual constituents against mosquitoes. *J Med Entomol* 50: 1267 - 1274.
- Bhattacharyya P, Chakrabarty PK, Chowdhury BK. 1985. Glycozolidol, an antibacterial carbazole alkaloid from *Glycosmis pentaphylla*. *Phytochemistry* 24: 882 - 883.
- Chaubey MK. 2007. Insecticidal activity of *Trachyspermum ammi* (Umbelliferae), *Anethum graveolens* (Umbelliferae) and *Nigella sativa* (Ranunculaceae) essential oils against stored-product beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Afr J Agr Res* 2: 596 - 600.
- Conti B, Leonardi M, Pistelli L, Profeti R, Ouerghemmi I, Benelli G. 2013. Larvicidal and repellent activity of essential oils from wild and cultivated *Ruta chalepensis* L. (Rutaceae) against *Aedes albopictus* Skuse (Diptera: Culicidae), an arbovirus vector. *Parasitol Res* 112: 991 - 999.
- Fisher DL. 1985. **Comparative life history studies of sexual and parthenogenetic *Liposcelis***, PhD Thesis, Reading University, UK.
- Fogang HPD, Womeni HM, Piombo G, Barouh N, Tapondjou LA. 2012. Bioefficacy of

- essential and vegetable oils of *Zanthoxylum xanthoxyloides* seeds against *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). **J Food Prot** 3: 547 - 555.
- Garcia M, Donael OJ, Ardanaz CE, Tonn CE, Sosa ME. 2005. Toxic and repellent effects of *Baccharis salicifolia* essential oil on *Tribolium castaneum*. **Pest Manag Sci** 61: 612 - 618.
- Govindachari TR, Pai BR, Subramaniam PS. 1966. Alkaloids of *Glycosmis pentaphylla* (Retz.) correa. **Tetrahedron** 22: 3245 - 3252.
- Greger H, Zehner G, Hofer O, Vajrodaya S. 1996. Bioactive amides from *Glycosmis* species. **J Nat Prod** 59: 1163 - 1168.
- Han QX, Chen JL, Liu BR, Tang HX. 2004. Toxic and repellent effects of several Chinese medicinal materials on the adults of *Tribolium castaneum* (Coleoptera: Tenebrionidae). **Plant Protection** 30: 60 - 63.
- Huang CJ. 1997. **Flora of China**. Science Press, Beijing, China.
- Huang YZ, Yang CJ, Xue D, Akinkulore RO, Yao YJ. 2007. Contact and repellency activities of ethanol extracts from 20 medicinal plants against *Rhizopertha dominica* (Fab.). **Acta Entomol Sin** 50: 118 - 124.
- Huang YZ, Lu YH, Li WJ. 2010. Repellent effect of the ethanol extracts from twelve herb plants on the adults of *Rhizopertha dominica* and *Tribolium castaneum*. **Chin Bull Entomol** 47: 387 - 392.
- Ito C, Itoigawa M, Sato A, Hasan CM, Rashid MA, Tokuda H, Mukainaka T, Nishino H, Furukawa H. 2004. Chemical constituents of *Glycosmis arborea*: three new carbazole alkaloids and their biological activity. **J Nat Prod** 67: 1488 - 1491.
- Khan MWY, Ahmad F, Ahmad I, Osman SM. 1983. Nonedible seed oils as insect repellent. **J Am Oil Chem Soc** 60: 949 - 950.
- Latha C, Joseph A. 1999. Evaluation of the larvicidal potential of *Glycosmis pentaphylla* against four important mosquito species of Kerala, India. **Int Pest Control** 41: 50 - 51.
- Liu ZL, Ho SH. 1999. Bioactivity of the essential oil extracted from *Evodia rutaecarpa* Hook f. et Thomas against the grain storage insects, *Sitophilus zeamais* Motsch. and *Tribolium castaneum* (Herbst). **J Stored Prod Res** 35: 317 - 328.
- Li ZH, Kucerova Z, Zhao S, Stejskal V, Opit G, Qin M. 2011. Morphological and molecular identification of three geographical populations of the storage pest *Liposcelis bostrychophila* (Psocoptera). **J Stored Prod Res** 47: 168 - 172.
- Lou CW, Sie MC, Lu CT, Kuo HL, Lin JH. 2013. A study of the effect of *Forcipomyia taiwana* repellents made of plant extracts. **Appl Mechanics Materials** 365 - 366: 1221 - 1224.
- Lukaseder B, Vajrodaya S, Hehenberger T, Seger C, Nagl M, Lutz-Kutschera G, Robien W, Greger H, Hofer O. 2009. Prenylated flavanones and flavanonols as chemical markers in *Glycosmis* species (Rutaceae). **Phytochemistry** 70: 1030 - 1037.
- Luo WR, Yang FD, Shi YB. 2012. Study progress of TCM curing methods. **Western J Trad Chin Med** 25: 108 - 112.
- Magan N, Hope R, Cairns V, Aldred D. 2003. Post-harvest fungal ecology: impact of fungal growth and mycotoxin accumulation in stored grain. **Eur J Plant Pathol** 109: 723 - 730.
- Mehmood F, Manzoor F, Khan ZD, Ali MI, Khan I, Rahim SMA. 2012. Evaluation of toxicity and repellency of essential oils of family Rutaceae against black ants (*Lasius niger*) in Pakistan. **Asian J Chem** 24: 3087 - 3090.
- Mills JT, Sinha RN, Demianyk CJ. 1992. Feeding and multiplication of a psocid *Liposcelis bostrychophilus* Badonnel (Psocoptera, Liposcelidae) on wheat, grain screenings and fungi. **J Econ Entomol** 85: 1453 - 1462.
- Nayak MK. 2006. **Psocod and mite pests of stored commodities: small but formidable enemies**. In I Lorini, B Bacaltchuk, H Beckel, D Deckers, E Sundfeld, JP dos Santos, JD Biagi, JC Celaro, LRDA Faroni, LOF Bortolini (eds.), Proceedings, 9th International Working Conference on Stored-Product Protection, Campinas, Sao Paulo, Brazil.
- Ngassoum MB, Tinkeu LSN, Ngatanko I, Tapondjou LA, Lognay G, Malaisse F, Hance T. 2007. Chemical composition, insecticidal effect and repellent activity of essential oils of three aromatic plants, alone and in combination, towards *Sitophilus oryzae* L. (Coleoptera: Curculionidae). **Nat Prod**

- Comm** 2: 1229 - 1232.
- Pandey AK, Singh P, Palni UT, Tripathi NN. 2013. Bioefficacy of plant essential oils against pulse beetles *Callosobruchus* spp. (Coleoptera: Bruchidae) in pigeon pea seeds with particular reference to *Clausena pentaphylla* (Roxb.) DC. **Arch Phytopathol Plant Prot** 46: 1408 - 1416.
- Phillips TW, Throne JE. 2010. Biorational approaches to managing stored-product insects. **Annu Rev Entomol** 55: 375 - 397.
- Quader MA, Nutan MTH, Rashid MA. 1999. Antitumor alkaloid from *Glycosmis pentaphylla*. **Fitoterapia** 70: 305 - 307.
- Sripisut T, Ritthiwigrom T, Promgool T, Yossathera K, Deachathai S, Phakhodee W, Cheenpracha S, Laphookhieo S. 2012. Glycopentaphyllone: The first isolation of hydroperoxyquinolone from the fruits of *Glycosmis pentaphylla*. **Phytochem Lett** 5: 379 - 381.
- Tang GW, Qiao LL, Su YP, Yang CJ, Cai WL, Hua HX. 2009a. Repellency and contact toxicity of extracts by different solvents from *Trigonella foenum-graecum* L. against two pest species. **J Chin Cereals Oils Assoc** 24: 102 - 108.
- Tang GW, Su YP, Qiao LL, Hua HX, Yang CJ. 2009b. The bioactivities of different extracts from *Trigonella foenum-graecum* against important stored-grain insect. **Grain storage** 38: 7 - 12.
- Tang GW, Yang WQ, Hu YJ, Yang CJ, Hua HX. 2009c. Bioactive effects of *Trigonella foenum-graecum* extracts by using different extract methods against *Tribolium confusum* Jacquelin du Val. **J Chin Cereals Oils Assoc** 24: 123 - 128.
- Throne JE, Opit GP, Flinn PW. 2006. **Seasonal distribution of psocids in stored wheat**. In I Lorini, B Bacaltchuk, H Beckel, D Deckers, E Sundfeld, JP dos Santos, JD Biagi, JC Celaro, LRDA Faroni, LOF Bortolini (eds.), Proceedings, 9<sup>th</sup> International Working Conference on Stored-Product Protection, Campinas, Sao Paulo, Brazil.
- Turner B. 1999. Psocids as pests: the global perspective. **Int Pest Control** 41: 185 - 186.
- Venkatachalam MR, Jebanesan A. 2001. Repellent activity of *Ferronia elephantum* Corr. (Rutaceae) leaf extract against *Aedes aegypti* (L.). **Bioresource Technol** 76: 287 - 288.
- Wang JS, Di YT, Yang XW, Li SL, Wang YH, Hao XJ. 2006. Hydroquinone diglycoside acyl esters from the stems of *Glycosmis pentaphylla*. **Phytochemistry** 67: 486 - 491.
- Wang JS, Yang XW, Di YT, Wang YH, Shen YM, Hao XJ. 2006. Isoflavone diglycosides from *Glycosmis pentaphylla*. **J Nat Prod** 69: 778 - 782.
- Wang MM. 2003. Introduction about several kinds of traditional Chinese medicine against storage maintenance experience. **Chin J Chin Mater Med** 28: 31.
- Wei LL, Hua RM, Li MY, Huang YZ, Li SG, He YJ, Shen ZH. 2014. Chemical composition and biological activity of star anise *Illicium verum* extracts against maize weevil, *Sitophilus zeamais* adults. **J Insect Sci** 14: 1 - 13.
- Xia CG, Chen JL, Li LS, Zheng YM, Yi JH. 2000. Repellent effect of moutan and its extract against several species of stored-product insects in Chinese medicinal materials. **Food Storage** 29: 3 - 9.
- Yang GZ, Wu Y, Chen Y. 2012. Alkaloids from the stems of *Glycosmis pentaphylla*. **Helv Chem Acta** 95: 1449 - 1454.
- Yao YJ, Yang CJ, Xue D, Huang YZ. 2007. Bioactivities of extracts from *Acorus gramineus* on four stored grain pests. **Acta Entomol Sin** 50: 309 - 312.
- Zettler JL, Arthur FH. 2000. Chemical control of stored product insects with fumigants and residual treatments. **Crop Prot** 19: 577 - 582.
- Zhou HY, Zhao NN, Du SS, Yang K, Wang CF, Liu ZL, Qiao YJ. 2012. Insecticidal activity of the essential oil of *Lonicera japonica* flower buds and its main constituent compounds against two grain storage insects. **J Med Plants Res** 6: 912 - 917.