

Isolation of α -spinasterol from *Amaranthus spinosus* stems

[Aislación de α -spinasterol de ramas de *Amaranthus spinosus*]

A.H.M Masum BILLAH¹, Mohammad M. HUSSAIN^{2*}, Mohammad G. DASTAGIR¹, Md. ISMAIL¹ & Abdul QUADER

¹Department of Chemistry, University of Dhaka, Dhaka - 1000, Bangladesh.

²Department of Pharmacy, Jagannath University, Dhaka - 1100, Bangladesh.

Contactos / Contacts: Mohammad M. HUSSAIN - E-mail address: m.musarrafhussain@gmail.com

Abstract

α -spinasterol was identified in stems of *Amaranthus spinosus*. The structure was obtained by spectroscopic methods and comparison with literature data.

Keywords: *Amaranthus spinosus*, Amaranthaceae, α - Spinasterol

Resumen

α -spinasterol fue identificado en ramas de *Amaranthus spinosus*. La estructura fue obtenida por métodos espectroscópicos y por comparación con datos de literatura.

Palabras Clave: *Amaranthus spinosus*, Amaranthaceae, α -Spinasterol .

Recibido | Received: May 3, 2011.

Aceptado en versión corregida | Accepted in revised form: May 26, 2012

Publicado en línea | Published online: January 30, 2013

Este artículo puede ser citado como / This article must be cited as: AHM Masum Billah, Mohammad M. Hussain, Mohammad G. Dastagir, Md. Ismail, Abdul Quader. 2013. Isolation of α -spinasterol from *Amaranthus spinosus* stems. *Bol Latinoam Caribe Plant Med Aromat* 12(1): 15 – 17.

INTRODUCTION

Amaranthus spinosus L. (Amaranthaceae) is an annual plant widely distributed in the humid zone of the tropics.

Diferent parts of the plan and extracts are used in traditional medicine. The seed are used as a poultice for broken bones (Duke *et al.*, 1985). The plant is astringent, diaphoretic, diuretic, emollient, febrifuge and galactogogue (Bown, 1995; Chopra *et al.*, 1986; Duke *et al.*, 1993). Methanolic extract of *Amaranthus spinosus* showed anti-inflammatory properties (Olajide *et al.*, 2004) and antimalarial activity (Hilou *et al.*, 2006)

Amaranthus grains and leaves are a good source of palmitic, oleic, and linoleic acids (Ghani, 2003; Kirtikar and Basu, 1980)

As part of a research project on bioactive compounds for the treatment of tropical diseases, isolated from medicinal plants, the preliminary phytochemical study of this plant growing on Bangladesh is reported.

MATERIALS AND METHODS

Plant Material

Representative samples of aerial parts of *Amaranthus spinosus* L. were collected from Chittagong District, Bangladesh. Voucher specimens (DACB accession number N35445) were deposited in the Bangladesh National Herbarium Dhaka, Bangladesh.


Extraction and Isolation

Air-dried and powdered stems of *A. spinosus* (125 g) were extracted in a soxhlet apparatus successively with petroleum ether (40 - 70°) and ethyl acetate. The extracts were filtered and concentrated, to yield the concentrated petroleum ether extract (13g) and ethyl acetate extract (10g).

Part of the petroleum ether extract (500 mg), was saponified with methanolic sodium hydroxide (0.5 M, 10 ml). Work up of the non-saponified material yield a white crystalline solid m.p. 151-154° C.

α -spinasterol (1). Solid, mp 151-154°; IR ν_{\max} (KBr): 3420(OH), 3050 (H-C=C), 2930, 3850, 1640, 1450, 1370, 1040, 970, 830. $^1\text{H-NMR}$ (CDCl_3) δ : 0.540 (s, 3H in C-18), 0.795 (broad s, 3H in C-27), 0.795 (3H in C-29), 0.814 (s, 3H in C-19), 0.847 (d, $J=5.9$ Hz in C-26), 1.024 (d, $J=6.67$ Hz in C-21), δ 1.40 – 2.0 (m, for methine and methylene protons), δ 2.009 (s, oxygenated methine proton at C-3), 3.584 (m, 1H at C-3), 5.024 (m, 1H, H-7), 5.045 – 5.119 (m, 2H, H-22 and H-23).

The compound was identified by direct comparison of the spectroscopic properties with previous published for α -spinasterol (1) (Agarwal, 1994-1995; Ismail *et al.*, 2010).



1

CONCLUSIONS

The chemical study of the stem of *Amaranthus spinosus* L. afforded one pure compound whose structure was established as α -spinasterol (1) by extensive spectroscopic studies as well as comparison with published data.

REFERENCES

- Agarwal OP. 1994-1995. **Chemistry of Organic Products**, 17th Edition, GOEL Publication House, Meerut, India.

- Bown D. 1995. **Encyclopaedia of Herbs and their Uses**. Dorling Kindersley, London, UK.
- Chopra RN, Nayar SL, and Chopra IC. 1986. **Glossary of Indian Medicinal Plants (Including the Supplement)**. Council of Scientific and Industrial Research, New Delhi, India.
- Duke JA, Ayensu ES. 1985. **Medicinal Plants of China**. Ref. Pub. Inc. Algonac., Michigan, USA.
- Finar IL. 1975. **Organic Chemistry**, Vol-2, Stereochemistry and the Chemistry of Natural Products, 5th edition, London, UK.
- Ghani, A. 1998. **Medicinal plants of Bangladesh: Chemical Constituents and Uses**. Asiatic Society of Bangladesh.
- He H, Corke H. 2003. Supercritical carbon dioxide extraction of oil and squalene from *Amaranthus* grain. **J Agric Food Chem** 51: 7921 - 7925.
- Hilou A, Nacoulma OG, Guiguemde TR. 2006. In vivo antimalarial activities of extracts from *Amaranthus spinosus* L. and *Boerhaavia erecta* L. in mice. **J Ethnopharmacol** 103: 236 - 240.
- Kirtikar KR, Basu BD 1980. **Indian Medicinal Plants**. Published by Singh B and Singh MP, India.
- Lamber JB, Herbert F. Shurvell, David LR Graham C. 1987. **Organic Chemistry**, Vol-2, Introduction to Organic Spectroscopy.
- Ismail M, Hussain MM, Dastagir MM, Billah M, Quader A, 2010. Phytochemical and antimicrobial investigation of *Luffa cylindrica*. **Bol Latinoam Caribe Plant Med Aromat** 9: 327 - 332.
- Olajide O, Ogunley B, Erinle T. 2004. Anti-inflammatory properties of *Amaranthus spinosus* leaf extract. **Pharmac Biol** 42: 521 - 525.
- Rastrelli L, Aquino R, Abdo S. 1998. Studies on the Constituents of *Amaranthus caudatus* Leaves: Isolation and Structure Elucidation of New Triterpenoid Saponins and Ionol-Derived Glycosides, **J Agric Food Chem** 46: 1797 - 1804.