

Artículo Original | Original Article

## Morphology, anatomy and histochemistry of the leaves of *Myracrodruon urundeuva* Allemão (Anacardiaceae)

[Morfología, anatomía e histoquímica de las hojas de *Myracrodruon urundeuva* Allemão (Anacardiaceae).]

Osmar Nascimento-SILVA<sup>1,2\*</sup>, Débora de Souza LEITE<sup>1</sup>, Luciana Aparecida BERNARDES<sup>1</sup>  
& Jose Geraldo Antunes de PAIVA<sup>1,2</sup>

<sup>1</sup>Laboratório de Botânica, Universidade Católica de Brasília, , QS 07-EPCT, 72.030-170, Águas Claras - Taguatinga, DF, Brazil

<sup>2</sup>Núcleo de Estudos Etnobotânicos e Etnofarmacognósticos - N.E.E.E da UCB

Contactos / Contacts: E-mail address: Osmar Nascimento-Silva [osmarns@gmail.com](mailto:osmarns@gmail.com)

### Abstract

The "aroeira" (*Myracrodruon urundeuva* Allemão) is a tropical tree with limited geographic distribution in South America, being found in drier formations such as the *Cerrado* and *Caatinga*. Empirically it is used with antiseptic, antiinflammatory, antiulcer, antidiarrhoeal and others. In this study we used mature leaves and expanded from the third and fourth nodes. Studies venation and morphology, anatomy and histochemistry were performed by the usual laboratory plant anatomy or the usual techniques of plant anatomy. For histochemical study of the fresh cuts various reagents and specific stains were used. The blade is elliptical leaflets with acute apex, oblique base, obtuse angle, entire margin and slightly wavy. Shows the pattern of venation feather-veined, pinnate type based generally asymmetrical and oblique. The indumentum is sericeous with trichomes deciduous. The consistency of the lamina is papyracea. The cuticle of leaflets, is thinner on the lower epidermis than on the upper epidermis, palisade parenchyma with a cell layer and spongy parenchyma with three cell layers with idioblasts containing crystals of CaCO<sub>3</sub>, tector trichomes simple multicellular with two or more cells are observed on both sides of the leaflets. The histochemical analysis revealed the presence of starch granules, crystals of calcium oxalate, fatty compounds, resins, phenolics and alkaloids compounds. The structural data obtained in this study may assist in ecophysiological characterization of the species and provide evidence for the identification of herbal medicines produced from that plant organ.

**Keywords:** *Myracrodruon urundeuva*; morphology; leaf anatomy; histochemistry.

### Resumen

El "aroeira" (*Myracrodruon urundeuva* Allemão) es una especie de árbol tropical con una distribución geográfica limitada en América del Sur, se encuentra en formaciones más secas, como el Cerrado y Caatinga. Empíricamente se usa en la cicatrización, como anti-inflamatorio y otros. En este estudio hemos utilizado las hojas maduras y ampliadas a partir de los nodos de tercero y cuarto. Para los estudios de venación y la morfología, la anatomía y los procedimientos histoquímicos fue el laboratorio de anatomía vegetal de costumbre. Los estudios de venación y morfología, anatomía e histoquímica fueron realizados por el laboratorio habitual de Anatomía vegetal o por las técnicas usuales de la anatomía vegetal. Los folíolos son elípticos con ápice agudo, base oblicua margen de ángulo obtuso todo ligeramente ondulado. El indumento es seríceo con tricomas de hoja caduca. La consistencia de la lámina es papirácea. Muestra el patrón de venación rectinervia, tipo pinnadas basan por lo general asimétrica y oblicua. La cutícula de los folíolos, es más delgada en la epidermis inferior de la epidermis superior, parénquima en empalizada con una capa de células y el parénquima esponjoso con tres capas de células idioblastos con drusas de oxalato de calcio, tricomas tectores multicelulares simples con dos o más células en la base se observado a ambos lados del folíolo. El análisis histoquímica reveló la presencia de gránulos de almidón, cristales de oxalato de calcio, compuestos grasos, resinas, fenoles y alcaloides. Los datos estructurales obtenidos en este estudio pueden ayudar en la caracterización ecofisiológica de la especie y aportar pruebas para la identificación de los medicamentos herbarios producidos a partir de ese órgano de la planta.

**Palabras Clave:** *Myracrodruon urundeuva*; morfología; anatomía de hojas; histoquímica

Recibido | Received: October 30, 2010.

Aceptado en versión corregida | Accepted in revised form: January 20, 2011.

Publicado en línea | Published online: January 30, 2011.

Este artículo puede ser citado como / This article must be cited as: Osmar Nascimento-SILVA, Débora de Souza LEITE, Luciana Aparecida BERNARDES and Jose Geraldo Antunes de PAIVA. 2011. Morphology, anatomy and histochemistry of the leaves of *Myracrodruon urundeuva* Allemão (Anacardiaceae). Bol Latinoam Caribe Plant Med Aromat 10(1): 56 – 66.

## INTRODUCTION

The family Anacardiaceae Lindl. is represented by approximately 70 genera and 700 species (Souza and Lorenzi, 2005), presenting predominantly pantropical distribution with some species in temperate regions (Cronquist, 1981). This family is characterized by the presence of secretory canals or ducts resiniferous latex and terpenes, also showing tannin compound and calcium oxalate crystals in parenchyma and silica in some cells of the xylem tissue (Engler, 1896, Metcalfe and Chalk, 1950; Cronquist, 1981).

*Myracrodruon urundeuva* Allemão, popularly known as "aroeira-do-sertão", is a tropical tree with geographic distribution, is restricted to South America, being found mainly in drier vegetation such as the "Cerrado" and the "Caatinga" (Rizzini, 1971; Lorenzi, 1992; Santin and Leitão-Filho, 1991; Florsheim, 1992; Pacheco et al., 2006). It features dense hard wood, with high concentration of tannins (Medina, 1966; Rizzini, 1971; Santos, 1987; Mainieri and Chimelo, 1989; Carvalho, 1994; Rizzini, 1995; Gonzaga et al. 2003), is included in the group of wood rot-proof timber (Nogueira, 1977). According to Medina (1966); Rizzini (1971), Nogueira (1977), Santos (1987); Mainieri and Chimelo, (1989); Carvalho (1994); Rizzini (1995) and Gonzaga et al. (2003) for its wood has all these qualities, is widely used mainly in rural areas in buildings such as fences and poles.

Together with other plant species with noble characteristics and high economic value, the "aroeira" has been widely exploited, leading to reduced size of natural populations, in many cases, extinction them completely (Brasil, 1992; Santos, 1993; Gonzaga et al., 2003; Freitas et al., 2005; Monteiro et al., 2010). Empirically, the "aroeira" is used to treat various illnesses, his bark is used as an antiseptic, healing, antiinflammatory, antiulcer, antidiarrhoeal, respiratory diseases and urinary tract (Matos, 1999; Gonzaga et al., 2003; Cabral and Carniello, 2004). Scientific studies have proven the anti-inflammatory, healing, antiulcer, anti-histamine, and analgesic antibradicinina, the shells of *M. urundeuva* (Viana et al., 1995; Rodrigues, 1999; Albuquerque et al. 2004). Other studies with out stem bark of *M. urundeuva* isolated tannin compound and chalcones - Urundeuvin A and B (Viana et al., 1995; Rodrigues, 1999; Albuquerque et al., 2004).

Moreover, a recent study Crivalero de Menezes et al. (2010), observed that the aqueous extract of *M. urundeuva* significantly reduced the oral

biofilm formation of *Streptococcus mutans* in rats, and the accumulation of *S. mutans* and enamel demineralization.

It is estimated that about 75% pure natural compounds used in the pharmaceutical industry were isolated following recommendations of folk medicine (Yunes et al., 2001). However, research to develop new drugs are time consuming and costly. In order to insure access of the poor to drug compounds, the WHO since 1976 encourages the use of medicinal plants as part of programs for primary health care (Calixto and Yunes, 2001; Cardoso and Verdecia, 1997), due to their efficacy combined with a low operating cost, resulting from the facility for the acquisition of plants. The use of this feature is very useful in the communities where the medical-pharmaceutical assistance is deficient, as is the case of the Brazilian Northeast, the region where mortality rates have remained above the national average (Matos, 1999).

It is known that the morpho-anatomical parameters of the leaves assist in the identification of pharmaceutical inputs and verifying the authenticity of drugs, avoiding possible adulteration, ensuring the proper use of them (Zanetti et al., 2004), also resulting in improved quality production of the species studied.

This study aims to understand the morphology and leaf anatomy, and thus contribute to a basic understanding and characterization of the venation pattern of *M. urundeuva*. Moreover, the results of histochemical tests serve as a basis for identifying classes of active biomolecules which can be used in the manufacture of herbal medicines more accessible mainly to the less favored population.

## MATERIALS AND METHODS

### Plant material

In this study expanded from the third and fourth nodes and mature leaves were used of *M. urundeuva*. The collection of plant material occurred in New Farm Nova Franca, Santa Maria da Vitória - BA in June 2007. After collection the leaves were kept at low temperature, and taken to the laboratory of Botany at the Catholic University of Brasília.

The species was previously identified through literature, comparison with material deposited in the herbarium of the Universidade de Brasília (UnB) and the ecological reserve of the Brazilian Institute of Geography and Statistics (IBGE) and later by experts in taxonomic groups. The herbarium specimens were collected according to the usual procedure and are

deposited in the herbarium of the Universidade Católica de Brasília.

### Morphologic studies

To study the venation, the leaves were cleared using techniques (Shobe and Lersten, 1967) with some modifications, with integral mounting of the leaves between two plates of glass, lacquer handicraft (Graciano-Ribeiro *et al.*, 2004, Paiva *et al.*, 2006). The venation patterns and morphological analysis followed Ash *et al.* (1999) and Vidal and Vidal (2003).

The organographic study was carried out observations with the naked eye and magnifying glass using LEICA MZ6. For the anatomical study, microscope OLYMPUS Cx31 was used.

### Anatomical studies

For the anatomical study transverse and longitudinal sections of median region and midrib of the leaflets were made which were made freehand with the aid of a cutting blade and a base of styrofoam. The sections were cleared in a solution of NaClO 2% according Shobe and Lersten (1967). The sections were washed in distilled water three times to remove the sodium hypochlorite. They were then stained with safranin/astra blue solution (Bukatsch, 1972).

To determine the type of stomatal apparatus were made in sections paradermic abaxial and adaxial surfaces with the aid of a cutting blade and forceps. The same sections were cleared in NaOH solution at 20% and 2% NaCl according Shobe and Lersten (1967), with some modifications. Thereafter, all the histological sections were mounted on semipermanent slides with glycerin gelatin Kaiser (Kaiser, 1880).

### Histochemical studies

For the histochemical study were done freehand cuts, the middle region and midrib of leaflets. the sections were subjected to various reactive dyes. the reagents used were: ethanolic sudan iii and sudan iv for detection of long-chain fatty such as lipids and other lipophilic compounds, and suberized walls cutinized (Foster, 1942); ferric chloride (Johansen 1940) and potassium dichromate (Gabe, 1968) for phenolic substances; lugol for the identification of starch grains (Sass, 1951); dilute sulfuric acid to test the nature of the crystals (Johansen, 1940); formalin with ferrous sulfate for evidence tannins (Schneider, 1977). For detection of alkaloids was used dragendorff reagent

(Yoder and Mahlberg, 1976), wagner's reagent and dittmar's reagent (Furr and Mahlberg, 1981).

The histochemical reactions were examined under a microscope OLYMPUS Cx31, and photographed with a digital camera attached to the eyepiece Sony DSC-P93. All photomicrographs were taken with digital camera Sony DSC-H1 coupled to said eyepiece microscope and magnifying glass. In all of the equipment ranges were obtained under the same conditions of optical images.

## RESULTS

### Leaflet and leaf morphology

*M. urundeuva* and has deciduous leaves are compound, petiolate, featuring a blade elliptic leaflets consistency papyraceous, acute apex, oblique base, obtuse angle, entire margin, with the absence of glands in limbo. The indumentum is sericeous, with multicellular trichomes that cover non-branched across the leaf surface. The average length of the leaf blade is 22.7 x 10.4 cm, and the leaflets 6.4 x 3.4 cm (Figure 1).


Shows the pattern of venation feather-veined, pinnate type. Venation of the second category is last with irregular marginal vein, the venation of the third category is dichotomized, the venation of the fourth category is regular polygonal reticulate. The veinlets are twice as white, the latter forming laciniosa entire marginal vein. Generally it has four ribs forming laciness (Figure 1).

### Leaflet anatomy


*M. urundeuva* presents a blade to leaflet anfi-hypostomatic. Paradermic sight the epidemics cells present sinusoids anticline walls. In the epidermis directed toward the adaxial face they are gifts stomata varying between anomocitic and tetracitic, surrounding the ribbings, being the anomocitic found in most frequency. In the abaxial face stomata of the anomocitic type are found (Figure 2).

In transversal section is observed thin cuticle thinn both the faces. The epidermis is unistratified, being the adaxial and abaxial face formed by tabular cells, with stomata in the same level of the usual epidemics cells (Figure 2). The limb to leaflet presents heterogeneous, collateral blade, mesophyl, where parenchyma palisade possess an only cellular layer and the apongy three cellular layers. Vascular bundles of the collateral type of small diameter are found in mesophyl. Idioblasts with crystals of oxalate of calcium in form of star, is found occasionally in parenchyma palisade (Figure 2).

**Figura 1.** Morphology and anatomy leaflets of *Myracrodruon urundeuva*. **a.** Morphology and general appearance of the leaf. **b.** Diaphanized leaflet, explaining the pattern of venation. **c, d** and **e.** Details of the venation pattern.



**Figura 2.** Sections transverse and paradermic of leaflets *Myracrodruon urundeuva*. **a.** Transverse section of middle third of the leaflets. **b.** Central vein. **c.** Leaflet stomata. **d.** Transversal section through leaflet midri. **e.** Transection of stomata on abaxial epidermis. **f.** Transection of stomata on adaxial epidermis. **g.** Non-glandular trichome. **h.** Glandular trichome. **i.** Aggregate crystals of calcium oxalate.



The main ribbing, in transversal section, presents a similar covering system with characteristics to the described ones for the remainder of the leaf. Below of the epidermis stratum of chollenchyma meets. Throughout the ribbing, is presented after lamellar parenchyma basic angular tending to lamellar in the abaxial and angular face in the adaxial face. The vascular system is formed by a vascular beam of the collateral type. Staple cellulosic fibers can be seen, next to the phloem and xylem. Involving the vascular beam an endoderm is observed. Secretory structures of the schizogenous type are found surrounding the vascular beam.

### Leaflet Histochemistry

Histochemical analysis showed that there is a greater impregnation of fatty compounds in the cuticle toward the adaxial side (Table 1). Phenolic compounds were found in almost every tissue leaflets in different concentrations (Table 1). They are found in abundance in the palisade and impregnating the periclinal walls of trichomes and the anticlinal and periclinal of walls cell collenchymatic and in smaller amounts in the cells surrounding the secretory structures (Table 1). Tannins were found in all tissues leaflets in large quantities (Table 1).

**Table 1.** Composite localization analyzed in leaflets of *Myracrodruon urundeuva* Allemão, by means of specific Histochemistry tests.

Tissue/anatomical structure	Phenolic composite		Alkaloids	Fatty compounds	Starch
	Generalities	Tannins			
Cuticle	-	-	-	+++	-
Epidermes	-	+	-	+	-
Trichome	++	++	-	++	-
Fundamental parenchyma	++	++	++	+	++
Palisade parenchyma	+++	+++	+	+	+
Spongy parenchyma	+	+++	+	+	+
Vascular bundles	+	++	++	+	-
Collenchyma	++	++	+	+	+

(+ + +) strong positive reaction for the composition, (++) positive reaction, (+) weakly positive reaction (-) negative reaction.

The alkaloids were present in moderate quantities in the fundamental parenchyma and pervading the bundle-sheath of parenchyma, and in smaller amounts in collenchymatic tissue and palisade and spongy mesophyll (Table 1). Starch granules were found in small amounts mainly in the fundamental parenchyma (Table 1).

### DISCUSIÓN

The morpho-anatomical data and histochemical are parameters that can be used in taxonomy and consequently assist in quality control of phytoterapics (Fank-de-Carvalho and Graciano-Ribeiro, 2005; Zanetti *et al.*, 2004).

The morphological data presented to elucidate the species morpho-anatomical studies relevant to the botanical family Anacardiaceae cited by Cronquist (1981), Ribeiro *et al.* (1999), Martinez-Millian and Cevallos-Ferriz (2005). Among them, anomocytic stomata, trichomes simple, crystals of calcium oxalate

and the presence of secretory structures in the midrib as well as studies conducted by Silva and Paiva (2007), which reveal that *Spondias tuberosa* Arruda has venation of the first and second category similar to those presented by *M. urundeuva*.

The presence of stomata on both sides of the blade leaflets may be related to environmental factors such as light intensity and relative humidity (Passos and Mendonca, 2006). According to Smith and colleagues (1997), this positioning of stomata reduces water loss by evapotranspiration. According to Pyykkö (1979) a greater abundance of stomata in abaxial prevent excess rainwater block the stomatal pore.

Transverse sections of leaves in *M. urundeuva* provide consistency to the analysis of the leaf with a thin cuticle, a list of environmental adaptation, since the characteristics leaflets are related to environmental characters. Silva and Paiva (2007), studying *S. tuberosa*, also observed a thin cuticle. Both *S. tuberosa* and *M. urundeuva* are in the same area soils and

phytogeographical, thereby warranting the similarity between species, since the environment is a determining factor for the phenotypic expression of the vegetative organs.

Idioblasts with prismatic (druse and/or raphide) in the form of crystals also occur along the vascular system. In this sense, once more the characteristic of the presence of these crystals in the venation of the Anacardiaceae is seen, as alleged by Martinez-Millan and Cevallos-Ferriz (2005) and Silva and Paiva (2007).

Leaflets of lamina mesophyll dorsiventral heterogeneous characteristics are also mesomorphic species has. This fact can be explained as a result of the specimens was collected in the forest transition in western Bahia. In this region rain 700-1800 mm per year concentrated in 83% of the wet season from October to April, with luminosity around 3,000 hours per year (Pinto *et al.*, 2006), features typical of mesophytic environment.

Vascular bundles of small caliber, which are found in *M. urundeuva*, can be interpreted as self-defense of the organism in the environment in which it is, since there are periods of drought, which would favor excessive loss of water in structures of thick bundles calibres. Such data can also be restated because the vascular bundle-being surrounded by parenchymatous sheath cells of light shot, which is nothing more than an endoderm cells that favors the xylem and phloem tissue of the water reserves in times of drought. Another relevant factor is the deciduousness. In general, when the drought is established, there is a drop leaf, allowing the plant, a survival without water loss. The presence of trichomes is another feature that provides the reflectance of light, avoiding high temperature inside leaflets, thus preventing the loss water (Valkama *et al.*, 2003). These trichomes are thick anticlinal cell walls and they act as mirrors, reflectors of sunlight, since the cuticle is deposited on the epidermal appendage, favoring the incidence of light rays.

The histochemistry analysis, showed presence of fatty compounds on epidermal tissue, this compound promotes water retention, and prevents excessive loss of this. The lignification of auxiliar cells of trichomes indicates the same stiffness, which makes their removal via mechanical difficult, favoring the fixing the same.

Phenolic compounds play an essential role in regulating plant growth and interaction with other agents, and promote plant chemical defense against

microorganisms, herbivores, pathogens, UV radiation, also possessing allelopathy (Croteau *et al.* 2000, Oliveira *et al.*, 2003, Taiz and Zeiger, 2004). From the standpoint of pharmacological activity they are antiseptic, anaesthetic, anti-inflammatory, antioxidant, have gonadotropic action, choleric, bile duct, antitumor, hipcholesterolemic, antipyretic and anti-flu (Diaz *et al.* 1999; Barbosa, 2004).

The alkaloids are considered the most important active compounds from the standpoint of pharmacology and medicine due to its physiological or psychological activity in humans, insects and other animals. They are often used as poisons, hallucinogens and stimulants (Salisbury and Ross, 1991). They have allelopathic action (Medeiros, 1990), antimicrobial, antifungal, and are toxic to some insects and mollusks (Robbers *et al.*, 1996). Yet such compounds to provide bitter taste plant organs (Kuklinski, 2000, Henriques *et al.*, 2002). According to Martins *et al* (1995), the alkaloids can be restricted to plant organs in different seasons of the year. According to Martin *et al* (1999), alkaloids have no definite action in plant organs, but Kuklinski (2000) cites a defensive action against pathogens, because they are toxic. Alkaloids have several pharmacological actions, some of them worth mentioning, as laxative, emetic, sedative for cough, antigout, antitumor (Cunha *et al.*, 2005a) antimalarial, antispasmodic (Cunha *et al.*, 2005b, Martin *et al.* 1999) stimulant, depressant central nervous system and hallucinogen (Cunha *et al.* 2005c; Kuklinski, 2000, Martin *et al.*, 1999).

Tannins are compounds found in greater abundance and in all plant tissues. Such compounds provide protection against dehydration plant, rotting (Macedo *et al.* 2005; Harbone, 1993; Von Teichman and van Wyk, 1994, Rocha *et al.*, 2002) and attack of pathogenic microorganisms (Scalbert, 1991; Trugilho, *et al.*, 2003). Another function related to these compounds is the protection against attack by herbivorous animals, because it blocks the action of digestive enzymes and can reduce fertility in moths (Souza and Marquete 2000). Pharmacologically, the tannins have astringent, healing, antiseptic, antioxidant (Kuklinski, 2000; Cunha and Batista, 2005), vasoconstrictors, haemostatic (Cunha and Batista, 2005) and antiinflammatory (Raphael and Kuttan, 2003; Osadebe and Okoye, 2003). It is believed that there are other functions related to them, although there is evidence that the work (Rocha *et al.*, 2002).

Tawaha *et al.* (2010) isolated new trimeric proanthocyanidin tannin, this compound was found to

have a potent inhibitory effect on COX-2, and exhibited moderate inhibition against COX-1.

Queiroz *et al.* (2002) found high amounts of phenolic compounds in the wood of *M. urundeuva*, about 20.2% of income earned for the crude methanol were gallic acid. Moreover, these authors tannins found in the wood are mainly the type of proanthocyanidins.

## CONCLUSIONES

The structural data obtained in this study may assist in ecophysiological characterization of the species and provide evidence for the identification of herbal medicines produced from that plant organ.

## ACKNOWLEDGEMENTS

The authors thank the Catholic University of Brasilia, in the person of Dra. Lourdes Loureiro, director of the biology course at UCB, to Dra. Cássia Beatriz Rodriguez Munhoz, head of the botany lab at UCB, by providing this space.

## REFERENCES

- Albuquerque RJM, Rodrigues LV, Viana GSB, 2004. Análise clínica e morfológica da conjuntivite alérgica induzida por ovalbumina e tratada com chalcona em cobaias. *Acta Cirúrgica Brasileira* 19: 43 - 68.
- Ash A, Ellis B, Hickey LJ, Johnson K, Wilf P, Wing S, 1999. Manual of leaf Architecture: Morphological description and categorization of dicotyledonous and net-veined monocotyledonous angiosperms. Leaf Architecture Working Group c/o Scott Wing. Department of Paleobiology. Smithsonian Institution. Washington – DC, Estados Unidos da América.
- Barbosa LCA, 2004. Introdução a Química Orgânica. Ed. Prentice Hall, São Paulo, p. 194 - 195.
- Brasil, Portaria no 006/92-N, de 15 de janeiro de 1992. Lista oficial de espécies da flora brasileira ameaçadas de extinção. Diário Oficial da República Federativa do Brasil, Brasília, DF, 23 jan. 1992.
- Bukatsch F. 1972. Bemerkungen zur Doppelfärbung Astrablau - Safranin. *Mikrokosmos* 61: 255.
- Cabral CDO, Carniello MA, 2004. Formas de uso medicinal da aroeira, *Myracrodruon urundeuva* Fr. All. em Porto Limão, Cáceres, MT. Trabalho apresentado no IV Simpósio sobre Recursos Naturais e Sócio-econômicos do Pantanal, realizado no período de 23 a 26 de Novembro de 2004, Corumbá-MS.
- Calixto JB, Yunes RA, 2001. Plantas medicinais sob a ótica da química medicinal moderna. Ed. Argos: Chapecó.
- Cardoso BAS, Porto Verdecia MP, 1997. Experiencia Cubana en el Estudio y Aplicación de Medicamentos Herbarios. *Revista Cubana de Plantas Medicinales* 2: 30 - 34
- Carvalho PER, 1994. Espécies florestais brasileiras: recomendações silviculturais, potencialidades e uso da madeira. Colombo: EMBRAPA-CNPQ; Brasília, DF: EMBRAPA-CNPQ, p. 672.
- Crivelaro de Menezes TE, Botazzo Delbem AC, Lourenção Brighenti F, Cláudia Okamoto A, Gaetti-Jardim E Jr. 2010. Protective efficacy of *Psidium cattleianum* and *Myracrodruon urundeuva* aqueous extracts against caries development in rats. *Pharm Biol* 48: 300 - 305.
- Cronquist A, 1981. An integrated system of classification of flowering plants. Columbia Univ. Press New York, Estados Unidos da América, p. 1262.
- Croteau R, Kutchan TM, Lewis N, 2000. Natural Products (Secondary Metabolites). *In: BUCHANAN, B., GRUISSEM, W. and JONES, R. Biochemistry and Molecular Biology of Plants. American Society of Plant Physiologists*, p. 1250 - 1318
- Cunha AP de, Batista MT, 2005a. Taninos. *In: CUNHA, A.P. de, 2005. Farmacognosia e Fitoquímica. Fundação Calouste Gulbenkian, Lisboa.*
- Cunha AP, Roque OR, 2005b. Compostos Fenólicos: Características e Origem Biossintética. p. 212 - 224. *In: Cunha, A. P., 2005. Farmacognosia e Fitoquímica. Fundação Calouste Gulbenkian, Lisboa, p. 670.*
- Cunha AP, Salgueiro L, Roque OR, 2005c. Alcalóides - aspectos gerais, p. 485-493. *In: Cunha, A. P., 2005. Farmacognosia e Fitoquímica. Fundação Calouste Gulbenkian, Lisboa, p. 670.*
- Díaz LB, Rodrigues TS, Giménez MDG, 1999. Ácidos Fenólicos y Fenoles Sencillos. Cumarinas y Lignanós, p. 191 - 206. *In: Fresno, A. M. Del, (editor), 1999. Farmacognosia General. Ed. Síntesis, Madri, p. 335.*
- Engler, A., 1896. Anacardiaceae: Anatomisches Verhalten. Leipzig: Die Natürlichen Pflanzenfamilien Bd.



- Fank de Carvalho SM, Graciano Ribeiro D, 2005. Arquitetura, anatomia e histoquímica das folhas de *Gomphrena arborescens* L.f. (Amaranthaceae). *Acta Botânica Brasílica* 19: 377 - 390.
- Florsheim SMB, 1992. Variações da estrutura anatômica e densidade básica da madeira de árvores de aroeira *Myracrodruon urundeuva* F. F. and M. F. Allemão (Anacardiaceae). Piracicaba - SP, 252p. Dissertação (Mestrado) - Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo.
- Foster, A.S. Practical plant anatomy. New York, Van Nostrand, 1942. 228p.
- Freitas MLM, Aukar APA, Sebbenn AM, Moraes MLT, Lemos EGM, 2005. Variabilidade genética intrapopulacional em *Myracrodruon urundeuva* Fr. All. por marcador AFLP. *Scientia Forestalis* 68: 21 - 28.
- Furr M, Mahlberg PG, 1981. Histochemical analyses of laticifers and glandular trichomes in *Cannabis sativa*. *J Nat Prod* 44: 153 - 159.
- Gabe M, 1968. Techniques histologiques. Masson and Cie, Paris.
- Gonzaga TWC, Mata MERMC, Silva H, Duarte MEM, 2003. Crioconservação de sementes de aroeira (*Astronium urundeuva* Engl.), E BARAÚNA (*Schinopsis brasiliensis* Engl.). *Rev Bras Prod Agroindustriais* 5: 145 - 154.
- Graciano Ribeiro D, Paiva JGA, Fank de Carvalho SM, Magalhães MP, 2004. Resina sintética brasileira uma alternativa econômica visando à substituição de resinas usuais. Proceedings of the 55<sup>o</sup> Congresso Nacional/26<sup>o</sup> Encontro Regional de Botânicos de MG, BA e ES, (Sociedade Botânica do Brasil, Viçosa, Minas Gerais, Brazil, 18 - 23 July) CD publication, no page number.
- Harbone JB, 1993. Ecological biochemistry. 4<sup>a</sup> ed. London: Academic Press, p. 1 - 318.
- Henriques ATH, Kerbe VA, Moreno PRH, 2002. Alcalóides: generalidades e aspectos básicos, p. 641-656. *In: Simões C. M. O., et al.* 2002. Farmacognosia da planta ao medicamento. 2<sup>a</sup> ed. Porto Alegre e Florianópolis: Editora da UFRGS e Editora da UFSC, p. 821.
- Johansen DA, 1940. Plant microtechnique. 3<sup>a</sup> ed. Paul B. Hoeber Inc. New York.
- Kaiser E, 1880. Verfahren zur Herstellung einer tadellosen Glycerin-Gelatine. *Botanisch Zentralbl* 180: 25 - 26.
- Kuklinski C, 2000. Farmacognosia: Estudio de las drogas y sustancias medicamentosas de origen natural. Ediciones Omega, Barcelona.
- Lima RJC, Moreno AJD, Castro SFL, Gonçalves JRS, Olivera AB, Sasaki JM, Freire PTC, 2006. Taninos Hidrolisáveis em *Bixa orellana* L. *Química Nova* 29: 507 - 509.
- Lorenzi H, 1992. Árvores brasileiras: manual de identificação e cultivo de plantas arbóreas nativas do Brasil. Nova Odessa: Plantarum.
- Macedo EG, Filho BGS, Potiguara RCV, Santos DSB, 2005. Anatomia e Arquitetura Foliar de *Montrichardia linifera* (Arruda) Schott (Araceae) Espécie da Várzea Amazônica. *Boletim do Museu Paraense Emílio Goeldi, Belém-PA* 1: 19 - 43.
- Mainieri C, Chimelo JP, 1989. Fichas características de madeiras brasileiras. São Paulo: IPT.
- Martinez Millan M, Cevallos Ferriz SRS, 2005. Arquitectura foliar de Anacardiaceae. *Rev. Mex Biodiversidad* 76: 137 - 190.
- Martín JJ, Moll MCN, Zurita AZ, 1999. Alcaloides, p. 251 - 262. *In: Fresno, A. M. Del, (editor),* 1999. Farmacognosia General. Ed. Síntesis, Madrid, p. 335.
- Martins ER, Castro DM, Castellani DC, Dias JE, 1995. Plantas medicinais. Ed. UFV, Viçosa-MG, p. 220.
- Matos FJA, 1999. Plantas de medicina popular do Nordeste: propriedades atribuídas e confirmadas. Fortaleza: Edições UFC.
- Medeiros ARM, 1990. Alelopatia: importância e suas aplicações. *Horti Sul* 1: 27 - 32.
- Medina JC, 1966. Flora do Brasil – dicotiledôneas arbóreas úteis. Instituto Agrônomo de Campinas 1-2: 925.
- Metcalf CR, Chalk L, 1950. Anatomy of dicotyledons. v. 1. Clarendon Press, Oxford.
- Monteiro JM, Lins Neto EM, Araújo ED, Amorim EL, Albuquerque UP, 2010. Bark regeneration and tannin content in *Myracrodruon urundeuva* Allemão after simulation of extractive damages-implications to management. *Environ Monit Assess* 10: 1770 - 1773.
- Oliveira RB, Godoy SAP, Costa FB. 2003. Plantas tóxicas: conhecimento e prevenção de acidentes. Ed. Holos, Ribeirão Preto-SP.
- Osadebe PO, Okoye EC, 2003. Anti-inflammatory effects of crude methanolic extract and fractions of *Alchornea cordifolia* leaves. *J Ethnopharmacol* 89: 19 - 24.

- Pacheco MV, Matos VP, Ferreira RLC, Feliciano AL, Pinto KMS, 2006. Efeito de temperaturas e substratos na germinação de sementes de *Myracrodruon urundeuva* Fr. All. (Anacardiaceae). *Revista Árvore* 30: 359 - 367.
- Paiva JGA de, Fank de Carvalho SM, Magalhaes MP, Graciano-Ribeiro D, 2006. Verniz vitral incolor 500: uma alternativa de meio de montagem economicamente viável. *Acta Botânica Brasílica* 20: 257 - 264.
- Passos MB, Mendonça MS, 2006. Epiderme dos segmentos foliares de *Mauritia flexuosa* L. f. (Arecaceae) em três fases de desenvolvimento. *Acta Amazônica* 36: 431 - 436.
- Pinto JM, Silva CL, Oliveira CS, 2006. Influência de variáveis climáticas e hidráulicas no desempenho da Irrigação de um pivô central no oeste baiano. *Revista Engenharia Agrícola* 26: 76 - 85.
- Pyykkö M, 1979. Morphology and anatomy of leaves from some woody plants in a humid tropical forest of Venezuelan Guyana. *Acta Botanica Fennica* 112: 1 - 41.
- Queiroz CRAA, Morais SAL, Nascimento EA, 2002. Caracterização dos taninos da aroeira-preta (*Myracrodruon urundeuva*). *Revista Árvore* 26: 485 - 492.
- Raphael KR, Kuttan R, 2003. Inhibition of experimental gastric lesion and inflammation by *Phyllanthus amarus* extract. *Journal Ethnopharmacol* 87: 193 - 197.
- Ribeiro JELS, Hopkins MJG, Vicentini A, Sothes CA, Costa MAS, Brito JM, Souza MAD, Martins LHP, Lohmann LG, Assunção PACL, Pereira EC, Silva CF, Mesquita MR, Procópio LC, 1999. Flora da Reserva de Ducke: Guia de identificação das plantas vasculares de uma floresta de terra-firme na Amazônia Central. Ed. INPA, Manaus, Brasil, p. 544-545.
- Rizzini CT, 1971. Árvores e madeiras úteis do Brasil: manual de dendrologia brasileira. São Paulo: Edgard Blücher.
- Rizzini CT, 1995. Árvores e madeiras úteis do Brasil - manual de dendrologia brasileira. 2ed. São Paulo: Edgard Blücher.
- Robbers EJ, Speedie KM, Tyler EV, 1996. Farmacognosia e Biotecnologia. Ed. Premier, São Paulo-SP, p. 372.
- Rocha JF, Rosa MMT, Frade CCM, Diersmann EM, 2002. Estudo Anatômico e Histoquímico em Folhas de *Plantago major* L. E *Plantago australis* Lam. (Plantaginaceae). *Revista Universidade Rural: Série Ciências da Vida* 22: 33 - 41.
- Rodrigues LV, 199. Análise morfológica e morfométrica da colite induzida por ácido acético, em ratos, e tratada com extratos vegetais (*Myracrodruon urundeuva* Fr. All.). 50f. Tese (Doutorado em Técnica Operatória e Cirurgia Experimental) - Universidade Federal de São Paulo, São Paulo, 1999.
- Salisbury FB, Ross CW, 1991. *Plant physiology*. 3ª. ed. California, Belmont: Wadsworth Publishing Company, p. 692.
- Santin DA, Leitão Filho HF, 1991. Restabelecimento e revisão botânica do gênero *Myracrodruon* Freire Allemão (Anacardiaceae). *Revista Brasileira de Botânica* 14: 133 - 145.
- Santos E, 1987. Nossas madeiras. Ed. Itatiaia, Belo Horizonte/MG.
- Santos GJC, 1993. Efeito biológico de *Trichoderma viride* Pers. ex S.F. Gray e *T. harzianum* Rifai no tratamento de sementes de aroeira do sertão (*Astronium urundeuva* (Fr. All.) Engl.) e na incorporação ao solo. Dissertação (Mestrado em Agronomia/Fitossanidade) - Escola Superior de Agricultura de Lavras, Lavras, MG.
- Sass JE, 1951. *Botanical Microtechnique*. 2ª ed. Iowa Press Building, Iowa.
- Scalbert A, 1991. Antimicrobial properties of tannins. *Phytochemistry* 30: 3875 - 3883.
- Schneider H, 1977. Indicator hosts for peace decline: Symptomatology, histopathology, and distribution of mycoplasma-like organisms in leaf veins. *Phytopathology* 65: 592 - 601.
- Shobe WR, Lersten NR, 1967. A technique for clearing gymnosperm leaves. *Botanical Gazette* 127: 150 - 152.
- Silva ON, Paiva JGA, 2007. Estudos morfológicos e anatômicos em folhas adultas de *Spondias tuberosa* Arruda (Anacardiaceae Lindley). *Boletim Latinoam Caribe Plant Med Aromat* 6: 36 - 43.
- Smith WK, Vogelmann TC, De Lucia EH, Bell DT, Shepherd KA, 1997. Leaf form and photosynthesis: Do leaf structure and orientation interact to regulate internal light and carbon dioxide? *BioScience* 47: 785 - 793.
- Souza VC, Lorenzi H, 2005. Botânica sistemática: guia ilustrado para identificação das famílias de angiospermas da flora brasileira, baseado *Boletim Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas*/65

- na APG II. Ed. Plantarum, Nova Odessa, Brasil.
- Souza RCOS, Marquete O, 2000. *Miconia tristis* Spring e *Miconia doriana* Cogn. (Melastomataceae): anatomia do eixo vegetativo e folhas. *Rodriguesia* 51: 133 - 142.
- Taiz L, Zeiger E, 2004. *Fisiologia vegetal*. Ed. Artmed, Porto Alegre, p. 309 - 334.
- Tawaha K, Sadi R, Qa'dan F, Matalka KZ, Nahrstedt A. 2010. A bioactive prodelphinidin from *Mangifera indica* leaf extract. *Z Naturforsch C*. 65: 322 - 326.
- Trugilho PF, Mori FA, Lima JT, Cardoso DP, 2003. Determinação do Teor de Taninos na Casca de *Eucalyptus* spp. *Cerne* 9: 246 - 254.
- Valkama E, Salminen JP, Koricheva J, Pihlaja K, 2003. Comparative analysis of leaves trichome structure and composition of epicuticular flavonoids in Finnish Birch species. *Annals of Botany* 6: 643 - 655.
- Viana GSB, Matos FJA, Bandeira MAM, Rao VS, 1995. Aroeira-do-sertão (*Myracrodruon urundeuva* Fr. All.): estudo botânico, farmacognóstico, químico e farmacológico, 2ª edição revisada e ampliada, Fortaleza, Edições UFC.
- Vidal WN, Vidal MRR, 2003. *Botânica - organografia: Quadros sinóticos ilustrados de fanerógamos*. Ed. UFV, Viçosa, Brasil.
- Von Teichman I, Van Wyk AE, 1994. The generic position of *Protorhus namaquensis* Sprague (Anacardiaceae): evidence from fruit structure. *Annals of Botany* 73: 175 - 184.
- Yoder LR, Mahlberg PG, 1976. Reactions of alkaloid and histochemical indicators in laticifers and specialized parenchyma cells of *Catharanthus roseus* (Apocynaceae). *American Journal of Botany* 63: 1167 - 1173.
- Yunes RA, Pedrosa RC, Cechinel Filho V, 2001. *Pharmaceutics and Phytotherapics: the need for development of the industry of phytopharmaceutics and phytotherapics in Brazil*. *Química Nova* 24: 147 - 152.
- Zanetti GD, Manfron MP, Hoelzel SCS, 2004. Análise morfo-anatômica de *Tropaeolum majus* L. (Tropaeolaceae). *Iheringia, Série Botânica* 59: 173 - 178.