

Artículo Original | Original Article

## Chemical composition and bioactivity of *Piper auritum* and *P. multiplinervium* essential oils against the red flour beetle, *Tribolium castaneum* (Herbst)

[Composición química y bioactividad de los aceites esenciales de *Piper auritum* y *P. multiplinervium* contra el escarabajo rojo de la harina, *Tribolium castaneum* (Herbst)]

Karina CABALLERO-GALLARDO<sup>1</sup>, Jesús OLIVERO-VERBEL<sup>1,2</sup>,  
Nayive PINO-BENÍTEZ<sup>2,3</sup> & Elena E. STASHENKO<sup>2</sup>

<sup>1</sup>*Environmental and Computational Chemistry Group. Campus of Zaragocilla.*

*Faculty of Pharmaceutical Sciences. University of Cartagena, Cartagena, Colombia.*

<sup>2</sup>*Chromatography Laboratory, Research Centre of Excellence, CENIVAM, Industrial University of Santander, Bucaramanga, Colombia.*

<sup>3</sup>*Natural Products Research Group. Technological University of Chocó, Quibdó, Colombia.*

Contactos / Contacts: Jesús OLIVERO-VERBEL - E-mail address: [joliverov@unicartagena.edu.co](mailto:joliverov@unicartagena.edu.co)

### Abstract

Stored grain insects have shown resistance to synthetic insecticides, fact that has promoted the use of vegetable species for integrated pest management. *Piper auritum* and *P. multiplinervium* are two plants from the Piperaceae family present in the department of Chocó, Colombia, one of the most important hot spots of biodiversity in the world. This study was conducted to determine the repellent activity and toxicity of essential oils (EOs) isolated from these plants against *Tribolium castaneum*, using the area preference and contact toxicity methods, respectively. *P. auritum* EO presented greater repellency than *P. multiplinervium*, the first showed 100% lethality at minimum tested exposure period (24 h) whereas the second reached 16% at 72 h. EOs were analyzed by gas chromatography-mass spectrometry. *P. auritum* major components were safrole (93.2%) and miristicine (4.3%), whereas for *P. multiplinervium* were  $\beta$ -elemene (9.0%), trans- $\beta$ -caryophyllene (5.3%) and caryophyllene oxide (4.1%). It is speculated that the repellent effect of *P. auritum* may be related to its safrole content, a known repellent. These results evidenced Piper species could be used for development of repellents against *T. castaneum*.

**Keywords:** Aromatic plants, repellent, toxicity, insects, bioactivity.

### Resumen

Los insectos de los granos almacenados han mostrado resistencia a los insecticidas sintéticos, hecho que ha promovido el uso de especies vegetales para el manejo integrado de plagas. *Piper auritum* y *P. multiplinervium* son dos plantas de la familia Piperaceae presentes en el departamento del Chocó, Colombia, uno de los puntos de biodiversidad más importantes del mundo. En este estudio fue determinada la actividad repelente y toxicidad de los aceites esenciales (AE) aislados de estas plantas contra *Tribolium castaneum*, utilizando el método de área de preferencia y toxicidad por contacto, respectivamente. El AE de *P. auritum* presentó mayor repelencia que el de *P. multiplinervium*, el primero mostró 100% de letalidad al menor tiempo de exposición (24 h), mientras que el segundo alcanzó el 16% a las 72 h. Los AEs fueron analizados por cromatografía de gases-espectrometría de masas. Los componentes principales de *P. auritum* fueron safrol (93.2%) y miristicina (4.3%), mientras que para *P. multiplinervium* fueron  $\beta$ -elemene (9.0%), trans- $\beta$ -cariofileno (5.3%) y óxido de cariofileno (4.1%). Se cree que el efecto repelente de *P. auritum* puede estar relacionado con su contenido de safrol, un repelente conocido. Estos resultados evidencian que las especies de Piper podrían ser utilizadas para el desarrollo de repelentes contra *T. castaneum*.

**Palabras Clave:** Plantas aromáticas, repelentes, toxicidad, insectos, bioactividad.

Recibido | Received: February 9, 2013

Aceptado en versión corregida | Accepted in revised form: April 15, 2013

Publicado en línea | Published online: January 30, 2014

**Declaración de intereses | Declaration of interests:** This work was supported by the University of Cartagena, Cartagena; CENIVAM, Bucaramanga; and Colciencias, Bogotá, Colombia. Grant RC-245-2011 Colciencias, Patrimonio Autónomo del Fondo Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación, Francisco José de Caldas, and the National Program for Doctoral Formation (COLCIENCIAS, 567-2012).

**Este artículo puede ser citado como / This article must be cited as:** K Caballero-Gallardo, J Olivero-Verbel, N Pino-Benitez, EE Stashenko. 2014. Chemical composition and bioactivity of *Piper auritum* and *P. multiplinervium* essential oils against the red flour beetle, *Tribolium castaneum* (Herbst). *Bol Latinoam Caribe Plant Med Aromat* 13(1): 10 - 19.

## INTRODUCTION

In a world where the use of synthetic chemicals is common in various areas of human life, and concern for sustainable product development has increased, natural resources emerge as a source of raw material solution for effective and environmentally friendly products. Nowadays, a number of synthetic insecticides and fumigants are used to protect grain storage. Infestation of storage grains is a very serious problem as various life stages of insects cause economic damage and deteriorates the quality of food grains and food products. These pests often cause extensive loss of products stored in tropical and semitropical environments (Isman, 2000). Several species of insect pests are known for attacking granaries and other food products since ancestral times. *Tribolium castaneum* is major pest food products (Mishra and Tripathi, 2011) the adult stage of *T. castaneum* is very active and can breed throughout the year in warm areas (Pugazhvendon *et al.*, 2009). They live two years or more, during which female produce nearly 1000 eggs (Shukla *et al.*, 2010). However, the use of synthetic pesticide for the management of these insect pests cause great hazards for environmental and toxic to non target livings (Mishra and Tripathi, 2011).

Over the years, pesticide use has led to a variety of problems such as environmental pollution, chemical residues in food grains, development of insecticide resistance and toxicity to non target organisms (Cosimi *et al.*, 2009; Sousa *et al.*, 2009). The increasing public awareness about pesticide safety and possible damage to the environment has resulted in greater attention being given to natural products for the control of food stored pests (Rajendran and Sriranjini, 2008). Among the group of natural molecules used to control insects, essential oils (EOs) area being considered as outstanding solutions for their effectiveness and versatility. In fact, volatility and chemical diversity makes them excellent repellents. The use EOs derived from aromatic plants as low-risk insecticides has increased considerably owing to their popularity with organic farmers and environmentally conscious consumers (Regnault-Roger *et al.*, 2012).

EOs have repellent, insecticidal, and growth-reducing effects on a variety of insects (Rajendran and Sriranjini, 2008; Ukeh *et al.*, 2009; Mondal and Khalequzzaman, 2010).

The demand for naturally active compounds such as EOs has move forward the necessity to search

these chemicals in hot spots for biodiversity. Colombia is a country with approximately 45.000 plant species, and it is among the first countries throughout Latin America and the world in terms of biodiversity heritage. However, little is known about its richened and the enormous possibilities for innovation and sustainable development from this natural resource (Stashenko *et al.*, 2010). At the same time, as people depend on synthetic repellents to control pests of stored products, EOs are exported without any added value, reducing the income for many families, and the ability to create jobs through the manufacture of products derived from these natural resources.

The bioactivity of a large number of essential oils and their constituents have been evaluated against *T. castaneum* (Jahromi *et al.*, 2012; Amin *et al.*, 2012; Zandi-Sohani *et al.*, 2013; Liang *et al.*, 2013). In Colombia, EOs from *Cananga odorata*, *Lippia alba*, *L. origanoides*, *Cymbopogon citratus*, *C. martinii*, *C. flexuosus*, *Eucalyptus citriodora*, and *Citrus sinensis* have been tested as repellents using this insect (Olivero-Verbel *et al.*, 2009, 2010; Caballero-Gallardo *et al.*, 2011, 2012). However, species from the Piperaceae family have not been studied. In this context, the aim of this paper was to evaluate the repellent and insecticidal properties of EOs isolated from *Piper auritum* and *P. multiplinervium* cultivated in Chocó, Colombia against *T. castaneum* Herbst, one of the most far-flung and destructive stored-product pests throughout the world (Zapata and Smagghe, 2010).

## MATERIALS AND METHODS

### *Plant Material*

Two plant species of the genus *Piper* were used in the assays, *P. auritum* and *P. multiplinervium* (Figure 1). Plant material was collected from Chocó, Colombia, at the municipalities of Quibdó and Lloró, respectively). The taxonomic identification of the plants was carried out in the Colombian National Herbarium (COL) of the Institute of Natural Sciences, Faculty of Sciences of the National University of Colombia (Bogotá) and stored with voucher specimens COL No. 512209 and COL No. 519977 for *Piper auritum* and *P. multiplinervium*, respectively.

Figure 1

*Piper auritum**Piper multiplinervium*

### Studied species from Chocó, Colombia.

#### *Extraction of essential oils*

Essential oils were extracted from fresh leaves by microwave-assisted hydrodistillation (MWHd) as previously described by Stashenko *et al.* (2004). The process used a Clevenger type hydrodistillation equipment placed inside a domestic microwave LG microwave MS- 1242 ZK model, set at 2450 MHz, 800 W. The extraction was conducted in 10 intervals; 10 min for each one with 5 min rest time each one. The oil was dried using anhydrous sodium sulphate.

#### *Gas chromatography–mass spectrometry (GC–MS)*

An aliquot of each essential oil (20  $\mu$ L) was dissolved in dichloromethane to a final volume of 1 mL, and then transferred to vials for GC–MS. The chromatographic analysis was performed using an Agilent Technologies 6890 Plus (Palo Alto, CA) GC coupled to an Agilent Technologies MSD 5975 selective detector mass equipped with a split/splitless injector port (1:50 split ratio), an automatic injector Agilent 7863, and a data system HP ChemStation. One microliter of solvent was injected into the GC–MS equipment for the corresponding chromatographic analysis. A 60m $\times$ 0.25mm i.d. $\times$ 0.25 $\mu$ m with 5% phenyl poly(methylsiloxane) stationary phase. The oven temperature was set at 45  $^{\circ}$ C for 5 min, then increased 4  $^{\circ}$ C/min up to 150 $^{\circ}$ C for 2 min, then to 5  $^{\circ}$ C/min up to 250  $^{\circ}$ C for 5 min, and finally at 10 $^{\circ}$ C/min up to 275 $^{\circ}$ C. Helium was used as a carrier gas with 16.47 psi column head pressure and 1 mL/min linear velocity. The components identification

were based on Kováts indices (Ik) and by comparison of the mass spectra fragmentation patterns with those found in databases or libraries (NIST02, Adams, Wiley7n) (Adam, 2004).

#### *Insects and rearing conditions*

All experiments were conducted in the laboratory using long-established colonies of *T. castaneum*. The strain was kept in glass containers covered with a plastic mesh. Insects were reared on a diet of whole oat flour and kept at 26  $\pm$  2  $^{\circ}$ C, with a relative humidity of 70 - 85%, and a 10:14 h light:dark photoperiod. Red flour beetle adults of both sexes were collected, and those with size between 3.0 and 4.5 mm, and 7-10 days old, were employed in the experiments (Caballero-Gallardo *et al.*, 2011).

#### *Repellency tests*

The repellent activity was measured using the area preference method (Taponjoui *et al.*, 2005; Olivero *et al.*, 2009; Olivero-Verbel *et al.*, 2010; Caballero-Gallardo *et al.*, 2011; Lü and Shi, 2012). The solutions of EOs were prepared in acetone, and in all cases, a volume of 0.5 mL was evenly applied to a half-filter paper disk to obtain the desired oil volume per area unit of 0.00002, 0.0002, 0.002, 0.02, and 0.2  $\mu$ L/cm<sup>2</sup>. The other half of the filter paper was treated with an equal volume of acetone as a vehicle control. Test areas consisted of 9 cm Albet DP 597125 filter paper cut in half (31.8 cm<sup>2</sup>). A formulation of IR3535 [ethyl 3-(N-acetyl-N-butylamino) propionate], a synthetic

repellent (Licciardi *et al.*, 2006; Faulde *et al.*, 2010) which has been approved in the USA for skin applications (World Health Organization, 2001), was employed as a positive control, utilizing the same experimental conditions as the oils. The treated and control half disks were air-dried for 10 min to remove the solvent, re-attached with adhesive tape, and kept in 90 mm glass Petri dishes. Twenty adults of *T. castaneum* of both sexes were released at the center of each filter paper disk. Dishes were covered and placed in darkness at  $26 \pm 2$  °C and relative humidity of 70–85%. The numbers of *T. castaneum* specimens on treated and untreated portions of the experimental paper halves were counted for each dish after 2 and 4 h exposure. Percentage repellency (PR) for a given treatment time was obtained using the formula:

$$\text{PR} = [(\text{Nc}-\text{Nt})/(\text{Nc}+\text{Nt})] \times 100$$

Where Nc and Nt are the number of insects on the untreated (control) and treated areas, respectively. Three replicates were used for each tested concentration of EO, and each assay was repeated twice.

#### Contact toxicity on filter papers

The contact toxicity on filter papers was conducted using filter paper discs (Albet DP 597125, 9 cm diameter) (Tapondjou *et al.*, 2005). Oils were dissolved in acetone at concentrations of 0.2, 0.5, 0.9 and 1.2  $\mu\text{L}/\text{cm}^2$ , and 1 mL of each solution was uniformly distributed on the surface of the paper that was then placed in glass Petri dishes. After 10 min, once the solvent had been evaporated, 20 unsexed adults were transferred into each disc and stored in darkness at  $26 \pm 2$  °C and  $75 \pm 10$  % RH (Olivero *et al.*, 2009; Olivero-Verbel *et al.*, 2010; Caballero-Gallardo *et al.*, 2011). Three replicates were employed for each treatment, repeating each assay twice. Mortality was recorded after 24, 48 and 72 h. IR3535 was utilized as a positive control. Insects were considered dead when no leg or antennal movements were recorded.

#### Data analysis

The paired t-test was utilized to compare mean number of insects on the treated and untreated area of the filter paper. Repellency or attractancy was established if significant differences occurred for positive or negative percentage repellency, respectively. Normal distribution and equality between variances were checked by Kolmogorov-Smirnov and Bartlett's tests,

respectively. Comparisons between mean PRs for evaluated EOs were obtained using ANOVA, with Dunn's post-test used to compare treated with control-vehicle groups. Statistical analysis was performed with GraphPad 3.00. The mean repellence concentration ( $\text{RC}_{50}$ ) values were calculated using probit analysis (Finney, 1971). For all purposes, significance was set at  $P < 0.05$ .

## RESULTS

The results of repellency assays for tested EOs are presented in Table 1. Data showed that at tested concentrations, both EOs were strongly repellent against *T. castaneum*. At the lowest assayed concentration (0.00002  $\mu\text{L}/\text{cm}^2$ ), the EOs isolated from *P. auritum* and *P. multiplinervium* showed attractant activity at both exposure times (PR:  $-35 \pm 9\%$ ,  $-30 \pm 17\%$  and  $-5 \pm 10\%$ ,  $-8 \pm 13\%$ , respectively). The commercial repellent IR3535 was less effective than the two tested EOs, when the organisms were exposed to 0.02 and 0.2  $\mu\text{L}/\text{cm}^2$ .

Significant differences were found for the PR between the essential oil of *P. auritum* and the commercial repellent (IR3535) at higher concentrations (0.02 and 0.2  $\mu\text{L}/\text{cm}^2$ ). However, for the essential oil isolated from *P. multiplinervium*, no such differences were recorded.

$\text{RC}_{50}$  values are shown in Table 2. Repellent action was highly dependent upon oil concentration and exposure time. *P. auritum* essential oil was the most active with a  $\text{RC}_{50}$  value of 0.002  $\mu\text{L}/\text{cm}^2$ .

The results for the contact lethality on filter paper for the examined EOs are shown in Table 3.

The EO from *P. auritum* had greater toxicity than that from *P. multiplinervium*. After 24 h exposure, the first EO caused  $81 \pm 2\%$  mortality at the lowest tested concentration (0.2  $\mu\text{L}/\text{cm}^2$ ), but at 0.9  $\mu\text{L}/\text{cm}^2$  196 this was 100%. However, for *P. multiplinervium*, the maximum tested concentration (1.2  $\mu\text{L}/\text{cm}^2$ ) and exposure time (72 h) only produced  $16 \pm 4\%$  lethality.

The results obtained to repellent and mortality of essential oils against *T. castaneum* were analyzed using Student's t-test. In the case of the repellent activity were no significant differences were observed between the two essential oils, whereas significant differences were observed to essential oil *P. auritum* (Table 3).

The IR3535 presented very little effect on the survival of *T. castaneum* adults. Chemical composition of EO isolated from the two *Piper* species, as resulted from GC-MS analysis, is presented in Table 4. The

results revealed quite different compositions between the two species. The essential oil of *P. auritum* was characterized by high phenylpropanoid content (97.5%), in which safrole (93.2%) was found to be the major constituent and miristicine the second (4.3%). Among monoterpene hydrocarbons,  $\gamma$ -terpinene was found at low concentration (0.3%). The *P. multiplinervium* oil presented a high content of sesquiterpene hydrocarbons (67%) with  $\beta$ -elemene (8.7%), trans- $\beta$ -caryophyllene (5.3%)  $\alpha$ -selinene (3.8%) and  $\beta$ -selinene (3.4%) as major components, as well as oxygenated sesquiterpenes (7.1%), with caryophyllene oxide (4.1%) and trans-nerolidol (3.0%) as representative compounds.

## DISCUSSION

Different plant products, especially EOs, have considerable potential as insecticides and repellents, and they are gaining tremendous importance for the management of stored products, as these mixtures are considered biodegradable and ecologically safe. The red flour beetle, *T. castaneum*, is one of the most widespread and pernicious destructive stored-product pests throughout the world (Zapata *et al.*, 2010; El-Sayed and Genan, 2012). Results presented here have shown that essential oils obtained from *Piper auritum* and *P. multiplinervium* possess good bioactivity toward this insect. In addition, these oils have similar or better repellent properties than the commercial repellent used as a positive control (IR3535).

Properties of EOs are directly related to their chemical composition (Ben Marzoug *et al.*, 2011), that in turn, are a function of the plant part extracted, time of collection and growth environment conditions, among others (Zapata and Smagghe, 2010). The number and concentration of chemicals present in an EO may vary dramatically, even within the same species (Zapata and Smagghe, 2010). In this study, *P. multiplinervium* and *P. auritum* had different biological activities, reflecting their distinct chemical nature. *P. multiplinervium* does not contain a major component representing at least 50% of the oil

composition, whereas *P. auritum* is mainly composed by safrole (93.2%). This chemically based difference may explain the acute and low-dose toxicity observed for *P. auritum* (100% mortality after 24 h exposure, and  $RC_{50} = 0.002 \mu\text{L}/\text{cm}^2$ ).

The bioactivity of the EOs results from interaction among structural components, particularly the major constituents, although, the other compounds in the oil may also have a vital function (Intirach *et al.*, 2012). In this study the principal compound of *P. auritum* was safrole (93.2%), a methylenedioxy compound (Maul *et al.*, 2011), reported to have toxic properties derived from its structure as benzene derivative (Tripathi *et al.*, 2009). The metabolism of this naturally occurring chemical leads to the formation of safrole 2',3'-oxide, an electrophilic compound that produces DNA adducts in vitro and in vivo (Shen *et al.*, 2012), probably explaining its reported ability to induce liver tumors in mice and rats (Rietjens *et al.*, 2005). Several studies have been published about safrole being the main component in several *Piper* species (Abreu *et al.*, 2002; Protti *et al.*, 2005; Sauter *et al.*, 2012; Souto *et al.*, 2012), and used alone, it has been reported to be effective against *T. castaneum* adults (Huang *et al.*, 1999).

This work has clearly revealed that *Piper* chemotypes can vary considerably in their composition and biological properties; and therefore, their presence in Colombian flora is an important potential for promissory bioactive molecules, in particular as repellents, as it has been also demonstrated for *P. aduncum*, a species from the Brazilian biodiversity that has been shown to work as a repellent against the ant *Solenopsis saevissima* (Smith) (Hymenoptera: Formicidae) (Souto *et al.*, 2012).

In short, reported data revealed, for the first time, the repellent and insecticidal activity of essential oils extracted from *P. auritum* and *P. multiplinervium* grown in the department of Chocó, Colombia, and this property could be related to the presence of some particular compounds.

Table 1

Essential oils	Time									
	2 Hours					4 Hours				
	Concentration ( $\mu\text{L}/\text{cm}^2$ )									
	0.00002	0.0002	0.002	0.02	0.2	0.00002	0.0002	0.002	0.02	0.2
<i>P. auritum</i>	$-35 \pm 9^{\text{ab}}$	$30 \pm 7^{\text{a}}$	$32 \pm 11^{\text{a}}$	$82 \pm 3^{\text{ab}}$	$90 \pm 4^{\text{ab}}$	$-30 \pm 17$	$38 \pm 10^{\text{a}}$	$45 \pm 9^{\text{a}}$	$55 \pm 12^{\text{a}}$	$87 \pm 3^{\text{a}}$
<i>P. multiplinervium</i>	$-5 \pm 10$	$7 \pm 13$	$42 \pm 11^{\text{a}}$	$70 \pm 8^{\text{a}}$	$82 \pm 6^{\text{a}}$	$-8 \pm 13$	$10 \pm 16$	$45 \pm 11^{\text{a}}$	$70 \pm 9^{\text{a}}$	$80 \pm 4^{\text{a}}$
IR3535 <sup>c</sup>	$7 \pm 13$	$17 \pm 9$	$37 \pm 15$	$50 \pm 4^{\text{a}}$	$72 \pm 4^{\text{a}}$	$5 \pm 14$	$12 \pm 8$	$42 \pm 15^{\text{a}}$	$50 \pm 7^{\text{a}}$	$72 \pm 6^{\text{a}}$

Percentage Repellency<sup>1</sup> (PR) after two exposure times for two essential oils against *T. castaneum*

<sup>1</sup>Values are mean (SE of six replicates).

<sup>a</sup>Significant difference between the number of the organisms on both the treated and untreated halves, using a paired t test ( $P < 0.05$ ).

<sup>b</sup>Significant difference between essential oils and the positive control (IR3535), using ANOVA, with Dunn's post-test.

<sup>c</sup>. Commercial repellent.

Table 2

RC<sub>50</sub> (95% Confidence intervals) values of the essential oils tested against *T. castaneum*

Essential oils	Exposure time (h)	RC <sub>50</sub> ( $\mu\text{L}/\text{cm}^2$ )	Regression parameters*	
			R	P
<i>P. auritum</i>	2	0.002 (0.001-0.005)	0.940	0.0602
	4	0.002 (0.001-0.008)	0.920	0.0798
<i>P. multiplinervium</i>	2	0.008 (0.004-0.016)	0.967	0.0334
	4	0.007 (0.003-0.015)	0.966	0.0336
IR3535 <sup>a</sup>	2	0.015 (0.005-0.048)	0.997	0.0002
	4	0.016 (0.006-0.045)	0.983	0.0027

\*. From probit analysis.

<sup>a</sup>. Commercial repellent.

Table 3

Essential oils	Concentration ( $\mu\text{L}/\text{cm}^2$ )	Mortality (%)		
		24 h	48 h	72 h
<i>Piper auritum</i>	0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0
	0.2	81 $\pm$ 2*	90 $\pm$ 4*	96 $\pm$ 2*
	0.5	92 $\pm$ 2*	98 $\pm$ 1*	99 $\pm$ 1*
	0.9	100 $\pm$ 0*	100 $\pm$ 0*	100 $\pm$ 0*
	1.2	100 $\pm$ 0*	100 $\pm$ 0*	100 $\pm$ 0*
<i>Piper multiplinervium</i>	0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0
	0.2	0 $\pm$ 0	3 $\pm$ 2	9 $\pm$ 4
	0.5	4 $\pm$ 2	7 $\pm$ 2	12 $\pm$ 1
	0.9	5 $\pm$ 2	8 $\pm$ 2	15 $\pm$ 5
	1.2	6 $\pm$ 2	10 $\pm$ 3	16 $\pm$ 4
IR3535*	0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0
	0.2	0 $\pm$ 0	0 $\pm$ 0	2 $\pm$ 1
	0.5	0 $\pm$ 0	0 $\pm$ 0	3 $\pm$ 1
	0.9	0 $\pm$ 0	1 $\pm$ 1	3 $\pm$ 1
	1.2	1 $\pm$ 1	2 $\pm$ 1	4 $\pm$ 2

Mortality rates in *T. castaneum* exposed to EOs and IR3535

\*. Significant difference between the two EOs, using Student's t-test ( $P < 0.05$ ).

<sup>a</sup>. Commercial repellent.

Table 4

KIDB5 <sup>a</sup>	Compound	Chemical Class	Relative Composition (%)	
			<i>Piper auritum</i>	<i>Piper multiplinervium</i>
1065	$\gamma$ -terpinene	Monoterpene hydrocarbons	0.3	
1290	Safrole	Phenylpropanoids	93.2	
1520	Miristicine		4.3	
1398	$\beta$ -elemene			8.7
1420	trans- $\beta$ -caryophyllene	Sesquiterpene hydrocarbons		5.3
1490	$\beta$ -selinene			3.4
1497	$\alpha$ -selinene			3.8
1566	trans-nerolidol	Oxygenated		3.0
1584	Caryophyllene oxide	sesquiterpenes		4.1

Chemical composition (%) of tested essential oils

<sup>a</sup>Kovats index on DB-5 column

## ACKNOWLEDGEMENTS

This work was supported by the University of Cartagena, Cartagena; CENIVAM, Bucaramanga; and Colciencias, Bogotá, Colombia. Grant RC-245-2011 Colciencias, Patrimonio Autónomo del Fondo Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación, Francisco José de Caldas, and the National Program for Doctoral Formation (COLCIENCIAS, 567-2012).

## REFERENCES

- Abreu AM, Sevegnani L, Machicado AR, Zimmermann D, Rebelo RA. 2002. *Piper mikanianum* (Kunth) steudel from Santa Catarina, Brazil-a new source of safrole. **J Essent Oil Res** 14: 361 - 363.
- Adams RP. 2004. **Identification of essential oil components by gas chromatography/quadrupole mass spectrometry**, 4th ed., Allured Publishing Corporation. Carol Stream, Illinois, USA.
- Amin R, Mondol R, Rahman F, Alam J, Habib R, Hossain T. 2012. Evaluation of insecticidal activity of three plant extracts against adult *Tribolium castaneum* (Herbst). **Biologija** 58: 37 - 41.
- Ben Marzoug HN, Romdhane M, Lebrihi A, Mathieu F, Couderc F, Abderraba M, Khouja M. L, Bouajila J. 2011. *Eucalyptus oleosa* essential oils: chemical composition and antimicrobial and antioxidant activities of the oils from different plant parts (stems, leaves, flowers and fruits). **Molecules** 16: 1695 - 1709.
- Caballero-Gallardo K, Olivero-Verbel J, Stashenko E. 2011. Repellent activity of essential oils and some of their individual constituents against *Tribolium castaneum* Herbst. **J Agric Food Chem** 59: 1690 - 1696.
- Huang Y, Ho SH, Kini RM. 1999. Bioactivities of Safrole and Isosafrole on *Sitophilus zeamais* (Coleoptera: Curculionidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae). **J Econ Entomol** 92: 676 - 683.
- Caballero-Gallardo K, Olivero-Verbel J, Stashenko E. 2012. Repellency and toxicity of essential oils from *Cymbopogon martinii*, *Cymbopogon flexuosus* and *Lippia origanoides* cultivated in Colombia against *Tribolium castaneum*. **J Stored Prod Res** 50: 62 - 65.
- Cosimi S, Rossi E, Cioni PL, Canale A. 2009. Bioactivity and qualitative analysis of some essential oils from Mediterranean plants against stored-product pests: Evaluation of repellency against *Sitophilus zeamais* Motschulsky, *Cryptolestes ferrugineus* (Stephens) and *Tenebrio molitor* (L.). **J Stored Prod Res** 45: 125 - 132.
- El-Sayed HS, Genan MG. 2012. Food consumption and utilization by *Tribolium confusum* du Val (Coleoptera:Tenebrionidae) larvae and their susceptibility to the acetone extract of *Nerium oleander* L. (Apocynaceae) leaves in relation to three types of flour. **J Stored Prod Res** 51: 56 - 60.
- Faulde MK, Albiez G, Nehring O. 2010. Insecticidal, acaricidal and repellent effects of DEET- and IR3535-impregnated bed nets using a novel long-lasting polymer-coating technique. **Parasitol Res** 106: 957 - 965.
- Finney DJ. 1971. **Statistical Method in Biological Assay**, 2nd edn. Griffin, London, UK.
- Intirach J, Junkum A, Tuetun B, Choochote W, Chaithong U, Jitpakdi A, Riyong D, Champakaew D, Pitasawat B. 2012. Chemical constituents and combined larvicidal effects of selected essential oils against *Anopheles cracens* (Diptera: Culicidae). **Psyche** 2012: 1 - 11.
- Isman MB. 2000. Plant essential oils for pest and disease management. **Crop Prot** 19: 603 - 608.
- Jahromi MG, Pourmirza AA, Safaralizadeh MH. 2012. Repellent effect of sirinol (garlic emulsion) against *Lasioderma serricorne* (Coleoptera: Anobiidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae) by three laboratory methods. **African J Biotech** 11: 280 - 288.
- Liang Y, Li YL, Xu S, Zhao NN, Zhou L, Cheng J, Liu ZL. 2013. Evaluation of repellency of some chinese medicinal herbs essential oils against *Liposcelis bostrychophila* (Psocoptera: Liposcelidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae). **J Econ Entomol** 106: 513 - 519.
- Licciardi S, Herve JP, Darriet F, Hougard JM, Corbel V. 2006. Lethal and behavioural effects of three synthetic repellents (DEET, IR3535 and KBR 3023) on *Aedes aegypti* mosquitoes in laboratory assays. **Med Vet Entomol** 20: 288 - 293.

- Lü JH, Shi YL. 2012. The Bioactivity of essential oil from *Ailanthus altissima* swingle (Sapindales: Simaroubaceae) bark on *Lasioderma serricorne* (Fabricius) (Coleoptera: Anobiidae). **Adv Mat Res** 365: 428 - 432.
- Maul R, Siegl D, Kulling SE. 2011. Glucuronidation of the red clover isoflavone irilone by liver microsomes from different species and human UDP-glucuronosyl transferases. **Drug Metab Dispos** 39: 610 - 616.
- Mishra BB, Tripathi SP. 2011. Repellent activity of plant derived essential oils against *Sitophilous oryzae* (Linnaeus) and *Tribolium castaneum* (Herbst). Singapore. **J Sci Res** 1: 173 - 178.
- Mondal M, Khalequzzaman M. 2010. Toxicity of naturally occurring compounds of plant essential oil against *Tribolium castaneum* (Herbst). **J Biol Sci** 10: 10 - 17.
- Olivero J, Caballero K, Jaramillo B, Stashenko E. 2009. Actividad repelente de los aceites esenciales de *Lippia origanoides*, *Citrus sinensis*, y *Cymbopogon nardus* cultivadas en Colombia frente a *Tribolium castaneum*, Herbst. **Salud UIS** 41: 244 - 250.
- Olivero-Verbel J, Nerio L, Stashenko E. 2010. Bioactivity against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) of *Cymbopogon citratus* and *Eucalyptus citriodora* essential oils grown in Colombia. **Pest Manag Sci** 66: 664 - 668.
- Protti S, Fagnoni M, Albin A. 2005. Expedient synthesis of bioactive allylphenol constituents of the genus *Piper* through a metal-free photoallylation procedure. **Org Biomol Chem** 3: 2868 - 2871.
- Pugazhvendon SR, Elumalai K, Ross PR, Soundararajan M. 2009. Repellent activity of chosen plant species against *Tribolium castaneum*. **World J Zool** 4: 188 - 190.
- Rajendran S, Sriranjini V. 2008. Plant products as fumigants for stored-product insect control. **J Stored Prod Res** 44: 126 - 135.
- Regnault-Roger C, Vincent C, Arnason JT. 2012. Essential oils in insect control: Low-risk products in a high-stakes world. **Annu Rev Entomol** 57: 405 - 424.
- Rietjens IM, Boersma MG, van der Woude H, Jeurissen SM, Schutte ME, Alink GM. 2005. Flavonoids and alkenylbenzenes: mechanisms of mutagenic action and carcinogenic risk. **Mutat Res** 574: 124 - 138.
- Sauter IP, Rossa GE, Lucas AM, Cibulski SP, Roehe PM, Alves da Silva L, Brittes Rott M, 321 Figueiro Vargas RM, Cassel E, von Poser G. 2012. Chemical composition and amoebicidal activity of *Piper hispidinervum* (Piperaceae) essential oil. **Ind Crops Prod** 40: 292 - 295.
- Shen LC, Chiang SY, Lin MH, Chung WS, Wu KY. 2012. In vivo formation of N7-guanine DNA adduct by safrole 2',3'-oxide in mice. **Toxicol Lett** 18: 309 - 315.
- Shukla GS, Upadhyay VB, Mathur R, Prasad SG. 2010. Economic zoology, biostatistics and animal behaviour. Rustogi Publication, India, pp: 97.
- Sousa A, Faroni L, Pimentel M, Guedes R. 2009. Developmental and population growth rates of phosphine-resistant and susceptible populations of stored product insect pests. **J Stored Prod Res** 45: 241 - 246.
- Souto RNP, Harada AY, Andrade EHA, Maia JGS. 2012. Insecticidal activity of piper essential oils from the amazon against the fire Ant *Solenopsis saevissima* (Smith) (Hymenoptera: Formicidae). **Pest Manag** 41: 510 - 517.
- Stashenko E, Jaramillo B, Martinez J. 2004. Comparison of different extraction methods for the analysis of volatile secondary metabolites of *Lippia alba* (Mill.) N.E. Brown, grown in Colombia, and evaluation of its in vitro antioxidant activity. **J Chromatogr A** 1025: 93 - 103.
- Stashenko E, Martinez J, Ruiz C, Arias G, Duran C, Salgar W, Cala M. 2010. *Lippia origanoides* chemo type differentiation based on essential oil GC-MS and principal component analysis. **J Separ Sci** 33: 93 - 103.
- Tapondjou AL, Adler C, Fontem DA, Bouda H, Reichmuth C. 2005. Bioactivities of cymol and essential oils of *Cupressus sempervirens* and *Eucalyptus saligna* against *Sitophilus zeamais* Motschulsky and *Tribolium confusum* du Val. **J Stored Prod Res** 41: 91-102.
- Tripathi AK, Upadhyay S, Bhuiyan M, Bhattacharya PR. 2009. A review on prospects of essential oils as biopesticide in insect-pest management. **J Pharm Pharm** 1: 052 - 063.
- Ukeh DA, Birkett MA, Pickett JA, Bowman AS, Luntz AJM. 2009. Repellent activity of alligator pepper, *Aframomum melegueta*, and ginger, *Zingiber officinale*, against the maize weevil, *Sitophilus zeamais*. **J Phytochem** 70: 751 - 758.

- World Health Organization. 2001. **Report of the Fourth WHOPES Working Group Meeting.** WHO/CDS/WHOPES/2001.2. WHO, Geneva, Swizerland.
- Zandi-Sohani N, Hojjati M, Carbonell-Barrachina AA. 2013. Insecticidal and repellent activities of the essential oil of *Callistemon citrinus* (Myrtaceae) against *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). **Neotrop Entomol** 42: 89 - 94.
- Zapata N, Lognay G, Smagghe G. 2010. Bioactivity of essential oils from leaves and bark of *Laurelia sempervirens* and *Drimys winteri* against *Acyrtosiphon pisum*. **Pest Manag Sci** 66: 1324 - 1331.
- Zapata N, Smagghe G. 2010. Repellency and toxicity of essential oils from the leaves and bark of *Laurelia sempervirens* and *Drimys winteri* against *Tribolium castaneum*. **Ind Crop Prod** 32: 405 - 410.