



Artículo Original | Original Article

Ethnobotany and regional knowledge: combining popular knowledge with the biotechnological potential of plants in the Aldeia Velha community, Chapada dos Guimarães, Mato Grosso, Brazil

[Etnobotánica y conocimiento regional: Combinando conocimiento popular con el potencial biotecnológico de plantas en la comunidad de Aldeia Velha, Chapada dos Guimarães, Mato Grosso, Brasil]

Larissa Cavalheiro¹ & Germano Guarim-Neto²

¹*Instituto de Ciências Naturais, Humanas e Sociais, Universidade Federal de Mato Grosso, Campus de Sinop, Sinop/MT, Brasil*

²*Instituto de Biociências, Universidade Federal de Mato Grosso, Campus de Cuiabá, Cuiabá/MT, Brasil*

Contactos / Contacts: Larissa CAVALHEIRO - E-mail address: larissacavalheiro@gmail.com

Abstract: Ethnobotanical studies were conducted in the Aldeia Velha community (Mato Grosso, Brazil) through semi-structured interviews. The majority of participants were women who have resided for more than 20 years in the community. Seventy-two species (39 families) were cataloged; leaf tea was the most common form of preparation. Over 80% were medicinal, with much of it is used for inflammation and disorders of the respiratory, digestive or genitourinary systems. Finally, seven species occurring in the Chapada dos Guimarães cerrado have biotechnological potential, especially *Amburana cearensis*, *Mikania glomerata* and *Pseudobrickellia brasiliensis*, where conservation measures are necessary due to their high risk of extinction and/or vulnerability.

Keywords: Traditional knowledge, medicinal plants, regional biodiversity, plant conservation

Resumen: El presente trabajo muestra resultados de un estudio etnobotánico realizado en la comunidad Aldeia Velha (Mato Grosso, Brasil). La información de uso de las especies fue colectada a través de entrevistas semiestructuradas, aplicadas a mujeres que residen hace más de 20 años en ese local. Adicionalmente fueron catalogadas 72 especies distribuidas en 39 familias. Entre las especies catalogadas más del 80% fueron informadas como medicinales, generalmente utilizadas para el tratamiento de inflamaciones, trastornos del sistema respiratorio, digestivo y genitourinario. Entre las formas de preparación la infusión de hojas fue la forma de consumo más común. En este trabajo fueron identificadas especies presentes en el Cerrado Chapadense que muestran un importante potencial biotecnológico destacándose *Amburana cearensis*, *Mikania glomerata* y *Pseudobrickellia brasiliensis*; debido a su alto riesgo de extinción y/o vulnerabilidad por ello son necesarias medidas conservacionistas de esa biodiversidad.

Palabras clave: conocimiento tradicional, plantas medicinales, biodiversidad regional, conservación vegetal.

Recibido | Received: June 19, 2017

Aceptado | Accepted: December 22, 2017

Aceptado en versión corregida | Accepted in revised form: January 16, 2018

Publicado en línea | Published online: March 30, 2018

Declaración de intereses | Declaration of interests: To the Fundação de Amparo a Pesquisa do Estado de Mato Grosso (FAPEMAT) for the aid granted to the Doctoral Program in Biotechnology and Biodiversity of the Rede Pró-Centro-Oeste de Pós-graduação, Pesquisa e Inovação, and the Universidade Federal de Mato Grosso (UFMT) for the support offered.

Este artículo puede ser citado como / This article must be cited as: L Cavalheiro, G Guarim-Neto. 2018. Ethnobotany and regional knowledge: combining popular knowledge with the biotechnological potential of plants in the Aldeia Velha community, Chapada dos Guimarães, Mato Grosso, Brazil. *Bol Latinoam Caribe Plant Med Aromat* 17 (2): 197 - 216.

INTRODUCTION

Tropical biological diversity has been studied under three general questions: (i) description of the extraordinary diversity; (ii) understanding of its origins and maintenance; and (iii) functional estimates of how this diversity responds to natural and anthropogenic changes (Bawa *et al.*, 2004). In addition, we can include a fourth important function in the production of biotechnology, or bioproducts, from material such as wood, fibers, resins, organic chemicals and genes, expanding the knowledge for application in biotechnology, including medicines and cosmetics (Alho, 2008). This can help improve income distribution and sustainable regional development (Davidson & Artaxo, 2004). However, much of this biodiversity is under increasing threat from anthropogenic processes, primarily deforestation, agricultural activities, illegal logging (Morton *et al.*, 2006) and urban sprawl. Thus, it is probable that important components of biodiversity, as well as the environmental services they provide, are under some type of threat, and are neglected in the planning of environmental management, mainly due to a lack of information on distribution and ecology (Garrido-Filho, 2002; Feamside, 2006).

By using the knowledge of local populations acquired and passed down through the generations, as demonstrated by their interactions with the environment in which they live, allied and aligned with scientific research aimed at the sustainable management of tropical ecosystems, we can find sustainable solutions for the use of these ecosystems. In this regard, ethnobotany is a multidisciplinary science that unites and links anthropology and other disciplines with botany (Albuquerque, 1999), and provides a record of traditional botanical knowledge (Guarim-Neto *et al.*, 2000). This allows more extensive use of ecological information when exploiting plant resources (Prance, 1991), which can be used as a component of sustainable development in protected areas.

The plants are primarily used as a food source, although other uses are worth mentioning such as handicrafts, perfume, ritual and medicinal use. The popular use of medicinal plants is what many communities rely on as an alternative for treatment of diseases or maintaining health. However, their continuity can be threatened by the interference of factors external to the social

dynamics of the group, for example: the community's increased exposure to surrounding society and, consequently, external economic and cultural pressures (Amorozo & Gély, 1988; Amorozo, 2002); greater ease of access to modern medicine services (Nolan & Robbins, 1999; Lima *et al.*, 2000; Amorozo, 2002); and displacement of people from their natural environment to urban regions, which leads to the loss of the utilitarian character of popular knowledge accumulated over several generations and, consequently, to its disappearance (Valle, 2002).

In the vegetative context of Mato Grosso, the traditional and even contemporary knowledge of plants persists and diversifies in response to specific characteristics of the regions inhabited by communities, confirmed by numerous studies such as Farnsworth (1981), Guarim-Neto (1984), De La Cruz & Guarim-Neto (1996), Faria (1998), Pasa & Guarim-Neto (2005, 2006), Borba & Macedo (2006), Guarim-Neto & Carniello (2007) and Pasa (2007, 2011a, 2011b), among many others. Thus, Aldeia Velha in Chapada dos Guimarães show to this day their traditional knowledge of cerrado plants and even exotic cultivated species.

MATERIAL AND METHODS

Study area

The study area is located in the Baixada Cuiabana, in the Municipality of Chapada dos Guimarães, Mato Grosso (Brazil), about 60 km from the capital Cuiabá (Figure 1). It is delimited by the geographic coordinates 15°30' and 15°40' S and 55°35' and 55°50' W (MMA, 2009). The study's target community was the Aldeia Velha, also the name of the municipality's first district. The study area was chosen based on observations of traditional plant use within the Chapada dos Guimarães region, being used by traditional healers, witch doctors, herbal sellers and midwives – traditional personalities known within the region (Neves, 1994). According to Vieira Jr. *et al.* (2011), its vegetation is characterized by distinct phytophysionomies, with a prevalence of savannah grasslands with differing densities of trees and shrubs (*cerrado sentido restrito* and *campo sujo*). According to the records, Aldeia Velha (Figure 2) was the first settlement within the municipality.




Figure N° 1

Map of Mato Grosso and its biomes, with the study's target area – the Municipality of Chapada dos Guimarães highlighted

Botanical material collection

Botanical material indicated by the community was first identified by the popular name attributed to the plant by the participants, followed by the formal identification of the cited species with the scientific name according to the current classification system, APG IV (2016). Scientific names have been updated based on the *Flora do Brasil 2020* (2016), as well as the Plant List and the Missouri Botanical Garden (Tropicos) websites.

For plant specimens where there existed doubts about the nomenclature, 3 to 5 samples in good condition were collected for subsequent formal identification, i.e. identification keys and specialized literature were used to obtain the scientific name of the species. Specimens were dried, pressed and mounted. All material collected in a fertile state and of relative importance to the work was deposited at the *Centro-Norte-Mato-Grossense* Herbarium (CNMT), *Universidade Federal de Mato Grosso* (UFMT), Sinop campus, and was also deposited at the UFMT Central Herbarium, Cuiabá campus when requested.

Ethnobotanical survey

Interviews were carried out using semi-structured questionnaires to define the group's profile (Alexiades, 1996). Each participant completed and signed a consent form detailing the information to be passed on. The technique used for sampling and selection of participants was the "snowball" technique proposed by Bernard (1995) and described by Albuquerque & Lucena (2004), where from an initial contact, a "local specialist" interviewed indicates another and so on. This is one of the techniques used today, for two basic reasons: the timeliness and low cost involved in collecting the information (Maciel *et al.*, 2002; Albuquerque & Hanazaki, 2006). In addition to the recruitment of local experts, other participants who showed an interest in taking part in the research were also included. The research conducted was of an eminently qualitative nature. In total there were about 30 participants from a total of 17 residences, primarily from the oldest part.




Figure N° 2
Aldeia Velha community with collection points marked

For the purpose of data analysis, the medicinal applications of each plant were grouped and classified in accordance with the Economic Botany Data Collection Standard (EBDCS) described by Cook (1995) and presented in Table N° 1, and also used in the work of Bieski *et al.* (2015). The categories presented, including cultural disorders and rituals/magical uses, provide a picture that combines traditional medicine with ethnomedicine, expanding the range of medical/cultural disorders.

RESULTS AND DISCUSSION

Socioeconomic profile of the Aldeia Velha community

The participants were mostly married women ranging from 33 to 81 years of age. The number of children ranged from 1 to 12. The community primarily consisted of people who have been living there long-term (more than 20 years). Most participants had incomplete elementary schooling.

The results obtained support the majority of ethnobotanical works where older women (over 50 years) are the keepers of knowledge (Amorozo, 2002; Bieski *et al.*, 2012; Amorozo *et al.*, 2006; Amaral & Guarim-Neto, 2008; Nanyingi *et al.*, 2008; Giraldo & Hanazaki, 2010; Pasa & Ávila, 2010; Vásquez *et al.*, 2014; Bieski *et al.*, 2015; David & Pasa, 2015).

A worrying finding in this study is related to age group, where knowledge is eminently held by older individuals and young people do not seem to know or show interest in the subject. However, this result is expected because according to Baptistel *et al.* (2014), the older generation has greater experience and experimentation. According to Almeida *et al.* (2010) and Silva *et al.* (2011), the younger generations know less than the older ones, thus not representing a loss of knowledge or "acculturation." In contrast, some of the participants stated that they take the children with

them to collect plants and try to pass on that knowledge already acquired by their predecessors such as parents and grandparents. An interesting fact is the presence of a resident who has taken the

medicinal plants course offered by the Rural Union of the city, where the community's interest in the subject is maintained by these initiatives.

Table N° 1
Groups of cultural diseases according to Cook's Classification

Group	Systems of the body	Initials
I	Blood system disorder	SAG
II	Circulatory system disorder	CIR
III	Digestive system disorder	DIG
IV	Endocrine system disorder	END
V	Genitourinary system disorder	GEN
VI	Immune system disorder	IMU
VII	Infections and infestations	INF
VIII	Inflamation	IFM
IX	Injury	INJ
X	Mental disorder	MEN
XI	Metabolic system disorder	MET
XII	Musculoskeletal system disorder	MUS
XIII	Neoplasms	NEO
XIV	Nervous system disorder	NER
XV	Nutritional disorder	NUT
XVI	Pain	DOR
XVII	Poisoning	ENV
XVIII	Pregnancy and/or postnatal disorder	PAR
XIX	Respiratory system disorder	RES
XX	Skin, cellular and subcutaneous tissue disorder	DE
XXI	Unspecified medical disorders	NE

Data analyzed from the questionnaires was able to verify that 100% of the participants use the plants in their day to day life and cultivate them within their place of residence. Approximately 85% of respondents use the plants all the time (15% from time to time); 65% actively collect plants from their local woods or other places, those who do not actively seek out the plants have opportunistically collected, and 90% of respondents can recognize them.

Ethnobotanical survey

This study cataloged 72 species belonging to 39 families (Table N° 2). The families with the highest number of species recorded were: Asteraceae with nine species, Lamiaceae with seven species, Fabaceae with four species and

Poaceae, Rutaceae, Solanaceae and Zingiberaceae with three species each. In addition to possessing the highest number of species (Souza & Lorenzi, 2012), these families are also cosmopolitan, harboring both native cerrado and common exotic plants introduced into the country. According to Guarim-Neto & Morais (2003), the more common the plant, the greater the probability that it will be used by the population. For example Asteraceae is one of the largest botanical families; Fabaceae is one of the most representative families within the cerrado biome, while Lamiaceae, although smaller in number of species, is a popularly cultivated family due to its high aromatic potency and large number of exotic herb species commonly used by the population.

Table N° 2
List of plants recorded by residents and additional information

*Scientific name	Common name	Plant part utilized	Form of utilization	General indication
Adoxaceae				
<i>Sambucus nigra</i> L.	Sabugueiro	Leaf	Tea Bath	Measles, Chickenpox
Alismataceae				
<i>Echinodorus grandiflorus</i> (Cham. & Schltr.) Micheli	Chapéu-de-couro	Leaf	Tea	Inflammation, Diuretic
Amaranthaceae				
<i>Alternanthera brasiliana</i> (L.) Kuntze	Terramicina	Leaf	Tea Leaf juice in bandage	Inflammation, Wounds, Blood
<i>Dysphania ambrosioides</i> (L.) Mosyakin & Clemants	Erva-de-Santa-Maria, Mentrúz, Mastruz	Leaf	Tea Leaf juice in bandage	Vermifuge, Inflammation, Wounds
Amaryllidaceae				
<i>Allium sativum</i> L.	Alho	Garlic clove	<i>In natura</i>	Coughing, Seasoning
Apiaceae				
<i>Eryngium foetidum</i> L.	Coentro-do-Pará	Leaf	In natura	Seasoning
<i>Foeniculum vulgare</i> Mill.	Erva-doce	Fruit, seed	Tea, <i>In natura</i>	Stomach, Seasoning
Apocynaceae				
<i>Mandevilla velame</i> (A.St.-Hil.) Pichon	Velame	Root	Juice alcohol	Wounds
Areaceae				
<i>Acrocomia aculeata</i> (Jacq.) Lodd. ex Mart.	Chiclete-cuiabano Bocaiuva	Fruit	<i>In natura</i>	Food
<i>Mauritia flexuosa</i> L.f.	Buriti	Fruit, stem	Dried fruits, Bark Wood	Crafts
Asteraceae				
<i>Achyrocline satureioides</i> (Lam.) DC.	Marcela	Leaf	Tea	Stomach, Stomach pain
<i>Ageratum conyzoides</i> L.	Mentastro	Leaf	Tea	Colic in babies
<i>Artemisia absinthium</i> L.	Losna	Leaf	Tea	Vermifuge, Fever
<i>Baccharis crispa</i> Spreng.	Carqueja	Leaf	Tea	Stomach, Weight loss
<i>Galinsoga parviflora</i> Cav.	Picão-branco	Whole plant	Bath	Itch relief, Wounds
<i>Lactuca sativa</i> L.	Alface	Leaf	Tea, <i>In natura</i>	Soothing agent, Food
<i>Matricaria chamomilla</i> L.	Camomila	Leaf, flower	Tea	Stomach ache Soothing agent
<i>Mikania glomerata</i> Spreng.	Guaco	Leaf	Tea, Syrup	Cough, Influenza
<i>Pseudobrickellia brasiliensis</i> (Spreng.) R.M.King & H.Rob.	Arnica	Leaf	Juice alcohol	Inflammation
Bignoniaceae				
<i>Tabebuia aurea</i> (Silva Manso) Benth. & Hook.f. ex S.Moore	Paratudo	Leaf	Tea	Various

Bixaceae				
<i>Bixa orellana</i> L.	Colorau, Urucum	Leaf, Seed	Tea, Seeds in water	Diabetes, Blood
Convolvulaceae				
<i>Ipomoea carnea</i> Jacq.	Algodãozinho Algodão-roxo	Leaf	Tea, Leaf juice	Inflammation, Infection, Wounds, Colic, Uterus
Costaceae				
<i>Costus spicatus</i> (Jacq.) Sw.	Cana-do-brejo	Leaf, Rhizome	Tea	Kidney
Crassulaceae				
<i>Kalanchoe crenata</i> (Andrews) Haw.	Saião	Leaf	Leaf juice	Hernia, Lung, Insect bites
Cucurbitaceae				
<i>Lagenaria siceraria</i> (Molina) Standl.	Cabaça	Fruit	<i>In natura</i>	Craft
<i>Momordica charantia</i> L.	Melão-de-São- Caetano	Leaf, Fruit	Tea, Leaf juice	Dengue, Diabetes, Malaria, Food
Dioscoreaceae				
<i>Dioscorea alata</i> L.	Cará Cará-do-ar	Fruit Rhizome	<i>In natura</i>	Food
Equisetaceae				
<i>Equisetum giganteum</i> L.	Cavalinha	Whole plant	Tea	Rim, Bladder
Fabaceae				
<i>Amburana cearensis</i> (Allemão) A.C.Sm.	Imburana	Bark, Seed	In alcohol	Inflammation, Analgesic
<i>Poincianella pluviosa</i> var. <i>peltophoroides</i> (Benth.) L.P. Queiroz	Sibipiruna	Leaves	Tea	Inflammation
<i>Senna occidentalis</i> (L.) Link	Fedegoso	Leaf	Tea, Leaves in alcohol	Abortive, Fever
<i>Tamarindus indica</i> L.	Tamarindo	Fruit	<i>In natura</i>	Food
Gentianaceae				
<i>Centaurium erythraea</i> Rafn.	Tea-porrete Coração-de- Jesus	Leaf	Tea	Dengue, Malaria, Viral disease Stomach pain
<i>Tachia guianensis</i> Aubl.	Caferana	Leaf	Tea	Stomach
Lamiaceae				
<i>Melissa officinalis</i> L.	Erva-cidreira Melissa	Leaf Flowers	Tea Bath	Soothing agent Wounds
<i>Mentha pulegium</i> L.	Poejo	Leaf	Tea, Syrup	Cough, Influenza Colic in babies
<i>Mentha x piperita</i> L.	Hortelã	Leaf	Tea, <i>In natura</i>	Worms, Influenza Seasoning
<i>Ocimum basilicum</i> L.	Alfavaca	Leaf	Tea	Cough
<i>Plectranthus barbatus</i> Andr.	Boldo	Leaf	Tea, Leaf juice	Stomach, Liver
<i>Rosmarinus officinalis</i> L.	Alecrim	Leaf	Tea, <i>In natura</i>	Heart, Seasoning Soothing agent
<i>Salvia officinalis</i> L.	Sálvia	Leaf	<i>In natura</i>	Seasoning
Lythraceae				
<i>Lafoensia pacari</i> A.St.-Hil.	Mangaba-brava	Bark	Tea	Gastritis, Insect bites

Malvaceae				
<i>Guazuma ulmifolia</i> Lam.	Mutamba, Chico-magro	Stem/Vine	<i>In natura</i>	Crafts
<i>Malva sylvestris</i> L.	Malva	Leaf	Tea	Infection, Bronchitis, Ulcer
Moraceae				
<i>Ficus carica</i> L.	Figo	Leaf	Tea	Asthma, Bronchitis
<i>Morus nigra</i> L.	Amora	Leaf	Tea	Insomnia
Musaceae				
<i>Musa paradisiaca</i> L.	Banana	Pedúnculo floral (coração)	Tea	Cough
Myrtaceae				
<i>Eugenia uniflora</i> L.	Pitangueira	Leaf	Tea	Heart
<i>Psidium guajava</i> L.	Goiabeira	Bark, Shoots	Tea, <i>In natura</i>	Stomach pain Aphthous sores
Nyctaginaceae				
<i>Boerhavia diffusa</i> L.	Erva-tostão Pega-pinto	Leaf	Tea	Various
Pedaliaceae				
<i>Sesamum indicum</i> L.	Gergelim-preto	Seed	<i>In natura</i>	Food
Phyllanthaceae				
<i>Phyllanthus niruri</i> L.	Quebra-pedra	Leaf	Tea	Kidney
Phytolaccaceae				
<i>Petiveria alliacea</i> L.	Guiné	Leaf	Tea, Bath	Magic
Piperaceae				
<i>Piper anisum</i> (Spreng.) Angely	Jaborandi	Leaf	Tea	Fever, Cough
Plantaginaceae				
<i>Scoparia dulcis</i> L.	Vassourinha	Leaf, Roots	Juice	Wounds
Poaceae				
<i>Cymbopogon citratus</i> (DC.) Stapf	Capim-cidreira	Leaf	Tea	Soothing agent
<i>Cymbopogon winterianus</i> Jowitt ex Bor	Citronela	Leaf	In alcohol	Insect repellent
<i>Merostachys multiramea</i> Hack.	Taquá	Leaf	Bath	Wounds
Polygonaceae				
<i>Polygonum punctatum</i> Elliott	Erva-de-bicho	Leaf	Tea, Bath	Dengue, Fever, Cough
Rubiaceae				
<i>Morinda citrifolia</i> L.	Noni	Fruit	<i>In natura</i>	Cancer, Various
Rutaceae				
<i>Pilocarpus jaborandi</i> Holmes	Jaborandi Jaborandi-do- cerrado	Leaf	Tea	Headache, Hair, Kidney
<i>Ruta graveolens</i> L.	Arruda	Leaf	<i>In natura</i> , in alcohol	Headache, Insect repellent Magig
<i>Zanthoxylum rhoifolium</i> Lam.	Mamica-de- porca	Leaf, Roots, Bark	Tea	Stomach, Fever, Aphrodisiac

Solanaceae				
<i>Capsicum frutescens</i> L.	Pimenta	Fruit	<i>In natura</i>	Food
<i>Solanum paniculatum</i> L.	Jurubeba	Fruit	Syrup	Bronchitis
<i>Solanum sisymbriifolium</i> Lam.	Juá, Joá	Fruit	Juice	Boils
Urticaceae				
<i>Cecropia pachystachya</i> Trécul	Embaúba	Buds	Tea	Cough, Bronchitis
Verbenaceae				
<i>Stachytarpheta jamaicensis</i> (L.) Vahl	Gervão	Roots	Tea	Urinary
Xanthorrhoeaceae				
<i>Aloe vera</i> (L.) Burm.f.	Babosa	Leaf	Juice Tea	Hair, Hernia, Cancer, Prostate, Urine
Zingiberaceae				
<i>Alpinia zerumbet</i> (Pers.) B.L.Burtt & R.M.Sm.	Colônia	Leaf (latex)	Tea	Heart Cancer
<i>Curcuma longa</i> L.	Açafrão-da-terra	Rhizome	<i>In natura</i>	Influenza, Throat, Diabetes
<i>Zingiber officinale</i> Roscoe	Gengibre	Rhizome	Tea, <i>In natura</i>	Throat

***All scientific names were confirmed from data available from the *Flora do Brazil 2020*, The Plant List, and Tropicos.**

Plant forms represented from the study were 62.5% herbaceous, 22.23% trees, 9.72% shrubs and sub-shrubs and 5.55% were vines. The higher percentage of herbaceous plants was expected, as these are more common and also easier to collect.

Leaves were the most commonly used parts of the plant with 74.01% of respondents recording leaf use. This data is also in accordance with the literature, as leaves are more common (they are always present), easy to access (Guarim-Neto & Morais, 2003), safer to use, available all year (Albuquerque & Hanazaki, 2006) and it is in the leaves where most of the plant's secondary metabolites, or active ingredients, are concentrated (Gonçalves & Martins, 1998).

Among the forms of preparation for use of these plants, 70% of respondents used tea (infusion) as the most common preparation,

followed by the *in natura* form with 16.15%, leaf juice (maceration) with 9.23% and use in baths with 4.62%. This data corresponds with the works of Martins *et al.* (2000), Matos (2002), Reis & Bellini (2011), and Bieski *et al.* (2012, 2015) in which the most common form of preparation was tea (infusion); this method of heating the plant (usually leaves, flowers and other above ground parts) with water facilitates the extraction of the active ingredients, making the home remedy faster and more efficient (Castellani, 1999).

Within the sampled size, 81.94% were considered medicinal, 18.05% food and 4.17% used for handicrafts and rituals. Thus, to date, within the plants considered medicinal, the majority are used for inflammation and disorders of the respiratory, digestive, metabolic, and genitourinary systems, as can be observed in Figure N° 3.




Figure N° 3

Diseases cited by informants according to categories proposed by Cook, 1995 (see Table N° 1)

The results were in accordance with those expected in that the plants are primarily used for the treatment of more common diseases as well as primary care such as kidney problems, wounds, worm treatment, Influenza and stomach problems (Pasa *et al.*, 2005). In this study, one of the most cited plants was the Cotton or Purple Cotton (*Ipomoea carnea* Jacq.), very common in the sampled areas, with its application related to the treatment of inflammation and uterine infections as well as in the treatment of wounds (injuries). According to the literature, the use of latex and leaves (maceration) is reported with the presence of alkaloids with antibacterial, immunological and uterine treatment properties, mainly with animal

tests (Haraguchi *et al.*, 2003; Hueza *et al.*, 2003; Hueza *et al.*, 2005; Melendez & Capriles, 2006; Hueza *et al.*, 2007; Armien *et al.*, 2011).

Other plants commonly mentioned were Chamomile (*Matricaria chamomilla* L.), used as tea for its sedative effects and treatment of bellyache, Erva-de-Santa-Maria (*Dysphania ambrosioides* (L.) Mosyakin & Clemants) used as a worming agent, against inflammations and wounds, and Terramicina (*Alternanthera brasiliana* (L.) Kuntze), also used in the treatment of inflammation, wounds and blood problems; both used in the form of tea or macerated as patches.

Biotechnological potential of plants: some indications

From this survey seven plants drew attention as to the number of citations and their use by the

community. They are considered native to the Chapadense cerrado and their effectiveness has been proven in some related works, as can be observed in Table N° 3.

Table N° 3
Biotechnological potential of some Chapadense cerrado plants

Species	Occurrence	Description of uses and associated scientific works
<i>Amburana cearensis</i> (Allemão) A.C.Sm. (Fabaceae) – Imburana, Amburana-de-cheiro, Cerejeira-do-nordeste	Present in the phytogeographical domains of the Caatinga, Cerrado and Atlantic Forest. It has confirmed occurrence for the states of Alagoas, Bahia, Ceará, Paraíba, Pernambuco, Piauí, Goiás, Espírito Santo, Minas Gerais, Rio de Janeiro. In Mato Grosso few specimens were found in the chapadense cerrado.	MEDICINAL: common cold, bronchitis, asthma, anti-cancer properties, anti-inflammatory and muscle-relaxant, antibiotic, antimutagenic, influenza antimicrobial (Leal <i>et al.</i> , 2003; Marinho <i>et al.</i> , 2004; Canuto & Silveira 2006; Oliveira <i>et al.</i> , 2009; Figueredo <i>et al.</i> , 2013; Lima <i>et al.</i> , 2013; Lorenzi, 2014); AGRONOMICS: insect repellent, seed mutagen (Lima <i>et al.</i> , 2013; Lorenzi, 2014); ECONOMIC/COMMERCIAL: carpentry, perfumery (Lorenzi, 2014); ETHNOBOTANICAL: species vulnerability (Gomes <i>et al.</i> , 2008; Martinelli & Morais, 2013).
<i>Equisetum giganteum</i> L. (Equisetaceae) - Cavalinha	Typical plant of the cerrado and Atlantic forest. It occurs in the states of Rondônia, Bahia, Paraíba, Pernambuco, Piauí, Distrito Federal, Goiás, Mato Grosso, Mato Grosso do Sul, Espírito Santo, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Rio Grande do Sul e Santa Catarina.	MEDICINAL: diuretic, anti-inflammatory, healing, digestive, hypoglycemic, remineralizing, hypotensive and antioxidant, fungicide against stomatitis, diabetes (Caetano <i>et al.</i> , 2002; Teixeira & Melo, 2006; Vieira <i>et al.</i> , 2006; Agra <i>et al.</i> , 2008; Santo-Dantas <i>et al.</i> , 2008; Lopes-Correa, 2010; Reuter & Salvador-Garcia, 2010; Zeni & Bosio, 2011; Alvarce <i>et al.</i> , 2015); AGRONOMICS: protection against environmental stresses, pest attack, fungicidal action (Bertalot <i>et al.</i> , 2010; Mello & Budel, 2014); VETERINARY: antithymic activity (Alvin-Carneiro, 1948; Riet-Correa <i>et al.</i> , 2002).
<i>Lafoensia pacari</i> A.St.-Hil. (Lythraceae) – Mangava-brava, dedaleira, didal, pacari.	Typical plant of the cerrado. It occurs in the states of	MEDICINAL: gastritis, ulcer, antifungal action, antibiotic effect, antimicrobial

	Tocantins, Bahia, Maranhão, Distrito Federal, Goiás, Mato Grosso do Sul, Mato Grosso, Minas Gerais, São Paulo e Paraná.	activity, asthma (Solon <i>et al.</i> , 2000; Souza <i>et al.</i> , 2002; Guarim-Neto & Moraes, 2003; Lima <i>et al.</i> , 2006; Menezes <i>et al.</i> , 2006; Batista, 2008; Rogerio <i>et al.</i> , 2008; Cabral & Pasa, 2009; Silva-Jr. <i>et al.</i> , 2010; Sampaio <i>et al.</i> , 2011; Tamashiro-Filho <i>et al.</i> , 2012); VETERINARY: canine parasitosis (Rogerio <i>et al.</i> , 2003). ETHNOBOTANICAL: species vulnerability (Martinelli & Morais, 2013)
<i>Mikania glomerata</i> Spreng. (Asteraceae) - Guaco	Plant found in the Cerrado and Atlantic Forest biomes. It has a confirmed occurrence for the states of Bahia, Espírito Santo, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Rio Grande do Sul, Santa Catarina. Specimens were found in Mato Grosso growing in backyards, but there are reports of collection in the Chapadense cerrado.	MEDICINAL: anti-inflammatory, bronchodilator, ulcers, snake venom, antiparasitic activity, bactericidal, antiallergic, antidiarrheal (Frutuoso <i>et al.</i> , 1994; Fierro <i>et al.</i> , 1999; Valverde <i>et al.</i> , 2001; Maiorano <i>et al.</i> , 2005; Salgado <i>et al.</i> , 2005; Iscan <i>et al.</i> , 2006; Kwiecinski <i>et al.</i> , 2008; Napimoga & Yatsuda, 2010; Kwiecinski <i>et al.</i> , 2011). ETHNOBOTANICAL: species vulnerability (Martinelli & Morais, 2013).
<i>Momordica charantia</i> L. (Cucurbitaceae) – Melão-de-São-Caetano, Erva-de-São-Vicente, Fruta-de-cobra, melãozinho.	A climber of tropical climate originating in Asia and Africa and fully naturalized Brazil, mainly in the Amazon and Cerrado. It occurs in the states of Acre, Amapá, Tocantins, Bahia, Ceará, Maranhão, Paraíba, Pernambuco, Piauí, Rio Grande do Norte, Goiás, Mato Grosso do Sul, Mato Grosso, Minas Gerais, Rio de Janeiro, São Paulo, Rio Grande do Sul e Santa Catarina.	MEDICINAL: diabetes, anti-bacterial, antiulcerogenic, anti-inflammatory, antityldemic, hypoglycemic, antihypertensive, anticancer, abortive, larvicidal, immunosuppressive, abortive, antihelmintic, contraceptive, dysmenorrhoea, antimalarial, laxative, rheumatism, some cancers, dengue (Anila & Vijayalakshmi, 2000; Basch <i>et al.</i> , 2003; Grover & Yadav, 2004; Konishi <i>et al.</i> , 2004; Nerurkar <i>et al.</i> , 2006; Dans <i>et al.</i> , 2007; Fernandes <i>et al.</i> , 2007; Braca <i>et al.</i> , 2008; Kobori <i>et al.</i> , 2008; Tan <i>et al.</i> , 2008).
<i>Polygonum punctatum</i> Elliott (Polygonaceae) – Erva-de-bicho.	Common plant in the Amazônia, Caatinga, Cerrado, Mata Atlântica, Pampa, and Pantanal biomes. It occurs	MEDICINAL: decrease in vascular permeability, antipyretic action and decrease in blood pressure, fungicidal action; antimicrobial activity, anti-

	practically all over Brazil, in the states of Acre, Amazonas, Pará, Roraima, Alagoas, Bahia, Ceará, Maranhão, Paraíba, Pernambuco, Piauí, Sergipe, Distrito Federal, Goiás, Mato Grosso do Sul, Mato Grosso, Espírito Santo, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, Rio Grande do Sul, Santa Catarina.	inflammatory action, biological control against <i>Schistosoma mansoni</i> (Simões et al., 1989; Pontier et al., 1991; Gorzalczyk et al., 1996; Alves et al., 2001; Penna et al., 2001).
<i>Pseudobrickellia brasiliensis</i> (Spreng.) R.M.King & H.Rob. (Asteraceae) – Arnica, arnica-do-mato	Typical plant of the cerrado and very common in the Chapada dos Guimarães region. Distributed in the states of Amazonas, Tocantins, Rondônia, Piauí, Ceará, Bahia, Mato Grosso, Goiás, Distrito Federal, Mato Grosso do Sul, Minas Gerais e São Paulo.	MEDICINAL: anti-inflammatory activity, immunomodulatory, allelopathy (Amorim, 2012; Amorim et al., 2012, 2016; Sivieri et al., 2015; Ribeiro et al., 2016). ETHNOBOTANICAL: species vulnerability (Martinelli & Moraes, 2013).

Of the plants indicated as having biotechnological potential, almost half are included in some category of threat. This analysis is related to the extinction risk to which the species are subject to, with data obtained by consulting the Red Book of Flora (Martinelli & Moraes, 2013) and *Flora do Brazil 2020*. Species are divided into the threat categories vulnerable, critically endangered and endangered.

This large number of endangered species can be explained by the concept promoted by Diamond (1989), who attributed the extinction of species to four great pillars, called the "Evil Quartet": anthropogenic activities such as habitat destruction, over-exploitation, introduction of exotic species and extinction chains. More recently, the effects of climate change have emerged as a new potential threat (Thomas et al., 2004). Although synergistically all of these factors may result in a greater than expected risk (Brook et al., 2008). In Mato Grosso state, the conversion of natural habitats into areas of crops and pastures, as well as the expansion of the urban frontier, appear to be the main threats to biodiversity

conservation (Baillie et al., 2004).

Pseudobrickellia brasiliensis (Spreng.) R.M.King & H.Rob. (Arnica) and *Mikania glomerata* Spreng. (Guaco) are among the threatened plants. Another feature that can increase this number even further is related to the large number of endemic species such as *Mikania* that is among the 20 most diversified genera of Brazilian flora, with 70% of the national species considered endemic; of these, 40 were evaluated and more than 50% of them are threatened (Martinelli & Moraes, 2013).

Another plant cited by the participants and listed in the Red Book of Flora is *Amburana cearensis* (Allemão) A.C.Sm. (Fabaceae), considered vulnerable, but not in the main list. This species is framed in the list of species as possessing economic value with verified or projected decline. This is also one of those species that we indicate as having biotechnological potential. Fabaceae is a cosmopolitan family; however this species has a restricted distribution in the cerrado. *Lafoensia pacari* A.St.-Hil. (Lythraceae) is listed in the minor concern

category, but it is faced with risks due to exploitation and loss of habitat by deforestation and expansion of agricultural and urban borders.

CONCLUSIONS

Brazil is a megadiverse country, both in terms of biology and culture, and Mato Grosso, because it is situated within three large biomes (Cerrado, Pantanal and Amazonia), deserves special mention. Studies aimed at reporting on and conserving this biodiversity have been frequent and increasingly urgent. In Chapada dos Guimarães, the participants showed themselves to be aware of the importance of the natural resources offered by the regional cerrado, and seek to enjoy them positively, primarily in obtaining medicinal plants.

One of the concerns of Ethnobotany is related to the search for new plant derived products, especially for industrial application, such as new food plants, manufacturing and particularly in the discovery of new drugs derived from medicinal plants. In addition, there is concern with human development issues such as the maintenance of native populations, conservation of nature and the conscious use of resources, bringing scientific research closer to society.

The need for biodiversity conservation in the Chapada dos Guimarães region is becoming more urgent, particularly as this region possesses plant species that are under severe threat, whether through deforestation, urban sprawl and/or over-exploitation. Finally, seven species present within the Chapadense cerrado have biotechnological potential, particularly *Amburana cearensis*, *Mikania glomerata* and *Pseudobrickellia brasiliensis*, where conservation measures are deemed necessary due to the high danger of extinction and/or vulnerability.

ACKNOWLEDGEMENTS

To the *Fundação de Amparo a Pesquisa do Estado de Mato Grosso* (FAPEMAT) for the aid granted to the Doctoral Program in Biotechnology and Biodiversity of the *Rede Pró-Centro-Oeste de Pós-graduação, Pesquisa e Inovação*, and the *Universidade Federal de Mato Grosso* (UFMT) for the support offered.

REFERENCES

- Agra MF, Silva KN, Basílio IJLD, França PF, Barbosa-Filho JM 2008. Survey of medicinal plants used in the region Northeast of Brazil. **Braz J Pharmacogn** 18: 472 - 508.
- Alavarce RAS, Saldanha LL, Almeida NLM, Porto VC, Dokkedal AL, Lara VS. 2015. The beneficial effect of *Equisetum giganteum* L. against Candida biofilm formation: New approaches to denture stomatitis. **Evid Based Complement Alternat Med** doi: 10.1155/2015/939625
- Albuquerque UP. 1999. Manejo tradicional de plantas em regiões neotropicais. **Acta Bot Bras** 13: 307 - 315.
- Albuquerque UP, Lucena RFP. 2004. **Métodos e técnicas na pesquisa etnobotânica**. NUPEEA, Recife/PE, Brasil.
- Albuquerque UP, Hanazaki N. 2006. As pesquisas etnodirigidas na descoberta de novos fármacos de interesse médico e farmacêutico: fragilidades e perspectivas. **Braz J Pharmacogn** 16: 678 - 689.
- Alexiades MN. 1996. **Collecting ethnobotanical data: An introduction to basic concepts and techniques**. In: Alexiades MN (ed.) Selected guidelines for ethnobotanical research: a field manual. New York Botanical Garden, Nueva York, USA.
- Alho CJR. 2008. The value of biodiversity. **Braz J Biol** 68: 1115 - 1118.
- Almeida CFCBR, Ramos MA, Amorim ELC, Albuquerque UP. 2010. A comparison of knowledge about medicinal plants for three rural communities in the semi-arid region of northeast of Brazil. **J Ethnopharmacol** 127: 674 - 684.
- Alves TM, Ribeiro FL, Kloos H, Zani CL. 2001. Polygodial, the fungitoxic component from the brazilian medicinal plant *Polygonum punctatum*. **Mem Inst Oswaldo Cruz** 96: 831 - 833.
- Alvin-Carneiro P. 1948. Envenenamento por *Equisetum* sp. (Cavalinha). **Ceres** 8: 32 - 36.
- Amaral NC, Guarim-Neto G. 2008. Os quintais como espaço de conservação e cultivo de alimentos: um estudo na cidade de Rosário do Oeste (MG, Brasil). **Bol Mus Para Goeldi Ciênc Hum** 3: 329 - 341.

- Amorim MLL. 2012. **Estudo fitoquímico e ensaios biológicos de *Pseudobrickellia brasiliensis* (Spreng.) R.M. King & H. Rob (Asteraceae).** Dissertação de mestrado, Universidade Federal dos Vales do Jequitinhonha e Mucuri, Diamantina, Minas Gerais, Brasil.
- Amorim MLL, Ribeiro JM, Graef CFF. 2012. **Isolamento e identificação do ácido caurenóico em extrato hexânico de Hojas de *Pseudobrickellia brasiliensis*.** In: XXII Simpósio de Plantas Mediciniais, Bento Gonçalves, RS, Brasil.
- Amorim MLL, Godinho WM, Archanjo FC, Graef CFF. 2016. Chemical constituents of *Pseudobrickellia brasiliensis* (Spreng.) R.M.King & H.Rob. leaves (Asteraceae). **Rev Bras Plant Med** 18: 408 - 414.
- Amorozo MCM, Gély AL. 1988. Uso de plantas medicinais por caboclos do Baixo Amazonas, Barcarena, PA. **Bol Mus Para Goeldi Sér Botânica** 4: 47 - 131.
- Amorozo MCM. 2002. Uso e diversidade de plantas medicinais em Santo Antônio do Leverger, MT, Brasil. **Acta Bot Bras** 16: 189 - 203.
- Amorozo, MCM. 2006. **Agricultura tradicional, espaços de resistência e o prazer de plantar.** In: Albuquerque UP. (Org.). Atualidades em etnobiologia e etnoecologia. 2 ed. NUPEEA/Sociedade Brasileira de Etnobiologia e Etnoecologia, Recife/PE, Brasil.
- Anila L, Vijayalakshmi NR. 2000. Beneficial effects of flavonoids from *Sesamum indicum*, *Emblica officinalis* and *Momordica charantia*. **Phytother Res** 14: 592 - 595.
- APG IV. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. **Bot J Linn Soc** 181: 1 - 20.
- Armién AG, Tokarnia CH, Peixoto PVS, Barbosa JD, Frese K. 2011. Clinical and morphologic changes in ewes and fetuses poisoned by *Ipomoea carnea* subspecies *fistulosa*. **J Vet Diagn Invest** 23: 221 - 232.
- Baillie JEM, Hilton-Taylor C, Stuart S. 2004. **IUCN Red list of threatened species: A global species assessment.** IUCN, Gland, Switzerland and Cambridge, UK.
- Baptistel AC, Coutinho JMCP, Lins-Neto EMF, Monteiro JM. 2014. Plantas medicinais utilizadas na Comunidade Santo Antônio, Currais, Sul do Piauí: um enfoque etnobotânico. **Rev Bras Plant Med** 16: 406 - 425.
- Bash E, Gabardi S, Ulbricht C. 2003. Bitter melon (*Momordica charantia*): a review of efficacy and safety. **Am J Health Syst Pharm** 60: 356 - 359.
- Batista HL. 2008. **Atividade antimicrobiana de extratos vegetais de plantas do Estado de Tocantins.** Dissertação de mestrado, UFC, Ceará, Brasil.
- Bawa KS, Kress WJ, Nadkarni N, Lele S. 2004. Beyond paradise-meeting the challenges in Tropical Biology in the 21st Century. **Biotropica** 36: 437 - 446.
- Bernard HR. 1995. **Research methods in anthropology: qualitative and quantitative approaches.** Altamira Press, London, UK.
- Bertalot MJA, Carvalho-Pupatto JG, Rodrigues EM, Mendes RD, Buso D. 2010. **Controle alternativo de doenças no morango.** Associação Brasileira de Agricultura Biodinâmica. http://www.biodinamica.org.br/pdf/controlo_alternativo_de_doencas_no_morango.pdf
- Bieski IGC, Santos FR, Oliveira RM, Espinosa MM, Macedo M, Albuquerque UP, Martins DTO. 2012. Ethnopharmacology of medicinal Plants of the Pantanal Region (Mato Grosso, Brazil). **Evid Based Complement Alternat Med** <http://dx.doi.org/10.1155/2012/272749>
- Bieski IGC, Leonti M, Arnason JT, Ferrier J, Rapinski M, Violante IMP, Balogun SO, Pereira JFCA, Figueiredo RCF, Lopes CRAS, Silva DR, Pacini A, Albuquerque UP, Martins DTO. 2015. Ethnobotanical study of medicinal plants by population of Valley of Juruena Region, Legal Amazon, Mato Grosso, Brazil. **J Ethnopharmacol** 173: 383 - 423.
- Borba AM, Macedo M. 2006. Plantas medicinais usadas para a saúde bucal pela comunidade do Bairro Santa Cruz, Chapada dos Guimarães, MT, Brasil. **Acta**

- Bot Bras** 20: 771 - 782.
- Braca A, Siciliano T, D'Arrigo M, Germano MP. 2008. Chemical composition and antimicrobial activity of *Momordica charantia* seed essential oil. **Fitoterapia** 79: 123 - 125.
- Brook BW, Sodhi NS, Bradshaw CJA. 2008. Synergies among extinction drivers under global change. **Trends Ecol Evol** 23: 453 - 460.
- Cabral, PRF, Pasa MC. 2009. Mangava-brava: *Lafloensia pacari* A. St. -Hil. (Lythraceae) e a etnobotânica em Cuiabá, MT. **Rev Biodiversidade** 8: 2 - 21.
- Caetano N, Saraiva A, Pereira R, Carvalho D, Pimentel MCB, Maia MBS. 2002. Determinação de atividade antimicrobiana de extratos de plantas de uso popular como anti-inflamatório. **Rev Bras Farmacogn** 12: 132 - 135.
- Canuto KM, Silveira ER. 2006. Constituintes químicos Da Casca Do Caule De *Amburana cearensis* A.C. Smith. **Quim Nova** 29: 1241 - 1243.
- Castellani DC. 1999. **Plantas medicinais**. Agromídia software, Viçosa, MG, Brasil.
- Cook FEM. 1995. **Economic Botany Data Collection Standard. Prepared for the International Working Group on Taxonomic Databases for Plant Sciences (TDWG)**. Royal Botanic Gardens, Kew, UK.
- Dans AM, Villarruz MV, Jimeno CA, Javelosa MA, Chua J, Bautista R, Velez GG. 2007. The effect of *Momordica charantia* capsule preparation on glycemic control in type 2 diabetes mellitus needs further studies. **J Clin Epidemiol** 60: 554 - 559.
- David M, Pasa MC. 2015. As plantas medicinais e a etnobotânica em Várzea Grande, MT, Brasil. **Interações** 16: 97 - 108.
- Davidson EA, Artaxo P. 2004. Globally significant changes in biological processes of the Amazon Basin: results of the Large-scale Biosphere-Atmosphere Experiment. **Global Change Biol** 10: 519 - 529.
- De La Cruz MGM, Guarim-Neto G. 1996. **Plantas medicinais utilizadas por agentes de saúde em Cuiabá - MT: um estudo etnobotânico**. Anais do Simpósio de Plantas Medicinais do Brasil, 1996 (Simpósio de Plantas Medicinais do Brasil, Florianópolis/SC), Brasil.
- Diamond JM. 1989. **Overview of recent extinctions**. In: Western D, Pearl MC: Conservation for the Twenty-First Century. Oxford University Press, Oxford, UK.
- Faria APO. 1998. **O uso de plantas medicinais em Juscimeira e Rondonópolis, MT: um estudo etnoecológico**. Dissertação de Mestrado, Universidade Federal de Mato Grosso, Brasil.
- Farnsworth N. 1981. An approach utilizing information from traditional medicine to identify tumor-inhibiting plants. **J Ethnopharmacol** 3: 85 - 99.
- Fearnside PM. 2006. Desmatamento na Amazônia: dinâmica, impactos e controle. **Acta Amaz** 36: 395 - 400.
- Fernandes NP, Lagishetty CV, Panda VS, Naik SR. 2007. An experimental evaluation of the antidiabetic and antilipidemic properties of a standardized *Momordica charantia* fruit extract. **BMC Complement Altern Med** 7: 29.
- Fierro IM, Silva ACB, Lopes CS, Moura RS, Barjafigalho C. 1999. Studies on the anti-allergic activity of *Mikania glomerata*. **J Ethnopharmacol** 66: 19 - 24.
- Figueiredo FG, Ferreira EO, Lucena BFF, Torres CMG, Lucetti DL, Lucetti ECP, Silva JMFL, Santos FAV, Medeiros CR, Oliveira GMM, Colares AV, Costa JGM, Coutinho HDM, Menezes IRA, Silva JCF, Kerntopf MR, Figueiredo PRL, Matias EFF. 2013. Modulation of the Antibiotic Activity by Extracts from *Amburana cearensis* A. C. Smith and *Anadenanthera macrocarpa* (Benth.) Brenan. **BioMed Res Int** 2013: 1 - 5. <https://doi.org/10.1016/j.sjbs.2015.02.001>
- Flora do Brasil 2020 em construção. Jardim Botânico do Rio de Janeiro. <http://floradobrasil.jbrj.gov.br/>
- Fruitoso VS, Gurjão MRR, Cordeiro RSB, Martins MA. 1994. Analgesic and anti-ulcerogenic effects of a polar extract from leaves of *Vernonia condensata*. **Planta Med** 60: 21 - 25.
- Garrido Filho I. 2002. Manejo florestal: questões econômico-financeiras e ambientais.

- Estudos Avançados** 16: 91 - 106.
- Giraldi M, Hanazaki N. 2010. Uso e conhecimento tradicional de plantas medicinais no Sertão do Ribeirão, Florianópolis, SC, Brasil. **Acta Bot Bras** 24: 395 - 406.
- Gomes ECS, Barbosa J, Vilar FCR, Perez JO, Vilar RC, Freire JLO, Lima NA, Dias TJ. 2008. Plantas da caatinga de uso terapêutico: levantamento etnobotânico. **Engenharia Ambiental: Pesquisa e Tecnologia** 5: 974 - 985.
- Gonçalves MIA, Martins DTO. 1998. Plantas medicinais usadas pela população do município de Santo Antônio de Leverger, Mato Grosso, Brasil. **Braz J Pharm** 79: 56 - 61.
- Gorzalczany S, Acevedo C, Muschietti L, Martino V, Ferraro G. 1996. Search for antiinflammatory activity in Argentine Medicinal Plants. **Phytomedicine** 3: 181 - 184.
- Grover JK, Yadav SP. 2004. Pharmacological actions and potential uses of *Momordica charantia*: a review. **J Ethnopharmacol** 93: 123 - 132.
- Guarim-Neto G. 1984. Plantas utilizadas na medicina popular cuiabana - um estudo preliminar. **Rev UFMT** 4: 45 - 50.
- Guarim-Neto G, Santana SR, Silva JVB. 2000. Notas etnobotânicas de espécies de Sapiendaceae Jussieu. **Acta Bot Bras** 14: 327 - 334.
- Guarim-Neto G, Morais RG. 2003. Recursos medicinais de espécies do Cerrado de Mato Grosso: Um estudo Bibliográfico. **Acta Bot Bras** 17: 561 - 584.
- Guarim-Neto G, Carniello MA. 2007. **Etnoconocimiento e saber local: um olhar sobre poblaciones humanas e os recursos vegetais**. In: Albuquerque UP, Alves AGC, Araújo TAS (Orgs.). Povos e paisagens: etnobiologia, etnoecologia e biodiversidade no Brasil. NUPEEA/UFRPE, Recife/PE, Brasil.
- Haraguchi M, Gorniak SL, Ikeda K, Minami Y, Kato A, Watson AA, Nash RJ, Molyneux RJ, Asano N. 2003. Alkaloidal components in the poisonous plant, *Ipomoea carnea* (Convolvulaceae). **J Agric Food Chem** 51: 4995 - 5000.
- Hueza IM, Dagli ML, Gorniak SL, Paulino CA. 2003. Toxic effect of prenatal *Ipomoea carnea* administration to rats. **Vet Hum Toxicol** 45: 298 - 302.
- Hueza IM, Guerra JL, Haraguchi M, Asano N, Gorniak SL. 2005. The role of alkaloids in *Ipomoea carnea* toxicosis: A study in rats. **Exp Toxicol Pathol** 57: 53 - 58.
- Hueza IM, Guerra JL, Haraguchi M, Gardner DR, Asano N, Ikeda K, Gorniak SL. 2007. Assessment of the perinatal effects of maternal ingestion of *Ipomoea carnea* in rats. **Exp Toxicol Pathol** 58: 439 - 446.
- Iscan G, Kirimer N, Kürkçüoğlu M, Arabacı T, Küpeli E, Başer KHC. 2006. Biological activity and composition of the essential oil of *Achillea schischkinii* Sosn. and *Achillea aleppica* DC. sbsp. *aleppica*. **J Agric Food Chem** 54: 170 - 173.
- Kobori RF, Gioria R, Brunelli KR. 2008. **Impacto potencial das mudanças climáticas sobre as doenças do pimentão no Brasil**. In: Ghini R, Hamada E. (Orgs.) Mudanças climáticas - Impactos sobre doenças de plantas no Brasil, Embrapa Informação Tecnológica, Brasília/DF, Brasil.
- Konishi T, Satsu H, Hatsugai Y, Aizawa K, Inakuma T, Nagata S, Sakuda SH, Nagasawa H, Shimizu M. 2004. Inhibitory effect of a bitter melon extract on the P-glycoprotein activity in intestinal Caco-2-cells. **Braz J Pharmacol** 143: 379 - 387.
- Kwiecinski MR, Felipe KB, Schoenfelder T, Wiese LPL, Rossi MH, Gonzalez E, Felicio JA, Wilhelm-Filho D, Pedrosa RC. 2008. Study of the anti-tumor potential of *Bidens pilosa* (Asteraceae) used in Brazilian folk medicine. **J Ethnopharmacol** 117: 69 - 75.
- Kwiecinski MR, Felipe KB, Correia JFG, Ferreira EA, Rossi MH, Gatti FM, Wilhelm-Filho D, Pedrosa RC. 2011. Brazilian *Bidens pilosa* Linné yields fraction containing quercetin-derived flavonoid with free-radical scavenger activity and hepatoprotective effects. **Libian J Med** 6: 5651.
- Leal L.K., Nechio M, Silveira ER, Canuto KM, Fontenele JB, Ribeiro RA, Viana GS. 2003. Anti-inflammatory and Smooth Muscle relaxant activities of the hydroalcoholic extract and chemical

- constituents from *Amburana cearensis* A. C. Smith. **Phytother Res** 17: 335 - 340
- Lima RX, Silva SM, Kunivoshi YS, Silva LB. 2000. Etnobiologia de comunidades continentais da área de Proteção Ambiental de Guaraqueçaba, Paraná, Brasil. **Etnoecológica** 4: 33 - 55.
- Lima MRF, Ximenes ECPA, Luna JS, Sant'ana AEG. 2006. The antibiotic activity of some Brazilian medicinal plants. **Braz J Pharmacogn** 16: 300 - 306.
- Lima LR, Cavalcante RRL, Martins MCC, Parente DM, Cavalcante AAMC. 2013. Avaliação da atividade anti-dematogênica, antimicrobiana e mutagênica das sementes de *Amburana cearensis* (A. C. Smith) (Imburana-de-cheiro). **Rev Bras Plant Med** 15: 415 - 422.
- Lopes-Correa AC. 2010. **Avaliação da atividade anti-inflamatória da pomada e do extrato etanólico bruto da *Equisetum pyramidale* Goldn nas lesões cutâneas de ratos normais e diabéticos no processo de reparação tecidual.** Dissertação de mestrado, Universidade Federal do Mato Grosso do Sul, Brasil.
- Lorenzi H. 2014. **Árvores brasileiras: manual de identificação e cultivo de plantas arbóreas do Brasil.** Instituto Plantarum, Nova Odessa/SP, Brasil.
- Maciel MAM, Pinto AC, Veiga VE. 2002. Plantas medicinais: a necessidade de estudos multidisciplinares. **Quim Nova** 23: 429 - 438.
- Maiorano AV, Marcussi S, Daher MA, Oliveira CZ, Couto LB, Gomes OA, França SC, Soares AM, Pereira PS. 2005. Anti-phididic properties of the aqueous extract of *Mikania glomerata*. **J Ethnopharmacol** 102: 364 - 370.
- Marinho MG, Brito AG, Carvalho KA, Bezerra-Santos CR, Andrade LHC, Barbosa-Filho JM, Piuvezam MR. 2004. *Amburana cearensis* e *Cumarina* imunomodulam os níveis de anticorpos antígeno-específico em camundongos BALB/c Sensibilizados com ovalbumina. **Acta Farm Bonaerense** 23: 47 - 52.
- Martinelli G, Moraes MA. 2013. Livro vermelho da flora do Brasil. Instituto de Pesquisas do Jardim Botânico do Rio de Janeiro. <http://cncflora.jbrj.gov.br/LivroVermelho.pdf>
- Martins ER, Castro DM, Castellani DC, Dias JE. 2000. **Plantas Mediciniais.** Editora UFV, Viçosa/MG, Brasil.
- Matos FJA. 2002. **Farmácias Vivas: sistema de utilização de plantas medicinais projetado para pequenas comunidades.** Editora UFC, Fortaleza/CE, Brasil.
- Meléndez PA, Capriles VA. 2006. Antibacterial properties of tropical plants from Puerto Rico. **Phytomedicine** 13: 272 - 276.
- Mello M, Budel JM. 2014. *Equisetum* L. (Equisetaceae): Uma revisão. **Cadernos da Escola de Saúde** 9: 1 - 15.
- Menezes VM, Atallah NA, Lapa AJ, Catapani WR. 2006. Assessing the therapeutic use of *Lafoensia pacari* St. Hil. extract (mangava-brava) in the eradication of *Helicobacter pylori*: double-blind randomized clinical trial. **Helicobacter** 11: 188 - 195.
- Morton DC, Defries RS, Shimabukuro YE, Anderson LO, Arai E, Espírito-Santo FDB, Freitas R, Morissette J. 2006. Cropland expansion changes deforestation dynamics in the southern Brazilian Amazon. **Proc Natl Acad Sci Biol** 103: 14637 - 14641.
- Nanyingi MO, Mbaria JM, Lanyasunya AL, Wagate CG, Koros KB, Kaburia HF, Munenge RW, Ogara WO. 2008. Ethnopharmacological survey of Bamburu district, Kenya. **J Ethnobiol Ethnomed** 4: 14 - 26.
- Napimoga MH, Yatsuda R. 2010. Scientific evidence for *Mikania laevigata* and *Mikania glomerata* as a pharmacological tool. **J Pharm Pharmacol** 62: 809 - 820.
- Nerurkar PV, Lee YK, Linden EH, Lim S, Pearson L, Frank J, Nerurkar VR. 2006. Lipid lowering effects of *Momordica charantia* (bitter melon) in HIV-1-protease inhibitor-treated human hepatoma cells, HepG2. **Braz J Pharmacol** 148: 1156 - 1164.
- Neves JES. 1994. **Chapada dos Guimarães: da descoberta aos dias atuais.** Ed. UFMT, Cuiabá/MT, Brasil.
- Nolan JM, Robbins MC. 1999. Cultural

- conservation of medicinal plant use in the Ozarks. **J Hum Org** 58: 67 - 72.
- Oliveira AK, Diógenes FEP, Coelho MFB, Maia SSS. 2009. Alelopatia em extratos de Fruits de juazeiro (*Ziziphus joazeiro* Mart. - Rhamnaceae). **Acta Bot Bras** 23: 1186 - 1189.
- Pasa MC, Guarim-Neto G. 2005. Garrafadas medicinais de plantas de Mato Grosso (I): *Anemopaegma arvense* (Vell.) Steff. e *A. glaucum* Mart. ex DC. **Biodiversidade** 1: 28 - 38.
- Pasa MC, Soares JJ, Guarim-Neto G. 2005. Estudo etnobotânico na comunidade de Conceição-Açu (alto da bacia do rio Aricá-Açu, MT, Brasil). **Acta Bot Bras** 19: 195 - 207.
- Pasa MC, Guarim-Neto G. 2006. Plantas medicinais no Vale do Aricá, município de Cuiabá, Estado de Mato Grosso, Brasil. **Biodiversidade** 1: 10 - 27.
- Pasa MC. 2007. **Um olhar etnobotânico sobre as comunidades do Bambá, Cuiabá, MT.** Ed. Entrelinhas, Cuiabá/MT, Brasil.
- Pasa MC, Ávila G. 2010. Ribeirinhos e recursos vegetais: a etnobotânica em Rondonópolis, Mato Grosso, Brasil. **Interações** 11: 195 - 204.
- Pasa MC. 2011a. O saber local e a medicina popular: A etnobotânica em Cuiabá, Mato Grosso, Brasil. **Bol Mus Para Goeldi Ciênc Hum** 6: 179 - 196.
- Pasa MC. 2011b. Abordagem etnobotânica na Comunidade de Conceição-Açu, Mato Grosso, Brasil. **Polibotânica** 31: 169 - 197.
- Penna C, Marino S, Vivot E, Cruanes MC, Munoz JD, Ferraro G, Gutkind G, Martino V. 2001. Antimicrobial activity of Argentine plants used in the treatment of infectious diseases: isolation of active compounds from *Sebastiania brasiliensis*. **J Ethnopharmacol** 77: 37-40.
- Pointier JP, Théron A, Imbert-Establet D, Borel G. 1991. Erradication of a sylvatic focus of *Schistosoma mansoni* using biological control by competitor snails. **Biol Control** 1: 244 - 247.
- Prance GT. 1991. What is ethnobotany today? **J Ethnopharmacol** 32: 209 - 216.
- Reis SLA, Bellini M. 2011. Representações sociais: teoria, procedimentos metodológicos e educação ambiental. **Acta Sci-Hum Soc Scienc** 33: 149- 159.
- Reuter C, Salvador-Garcia C. 2010. Protocolo de fitoterapia na hipertensão arterial sistêmica dos municípios da BP3. <http://www.gege.agrarias.ufpr.br/plantastoxicas/arquivos/equisetum%20giganteum.pdf>
- Ribeiro JM, Amorim MLL, Rocha AS, Graef CFF, Nery MC. 2016. Atividade alelopática do extrato aquoso das Hojas de *Pseudobrickellia brasiliensis* sobre a germinação e crescimento inicial de alface e tomate. **Rev Cient Vozes dos Vales** 9: 1 - 11.
- Riet-Correa F, Riet-Correa GE, Schild A. 2002. Importância do exame clínico para o diagnóstico das enfermidades do sistema nervoso em ruminantes eqüídeos. **Pesq Vet Bras** 22: 161 - 168.
- Rogério AP, Fontanari C, Borducchi E, Keller AC, Rosso M, Soares EG, Albuquerque DA, Faccioli LH. 2008. Anti-inflammatory effects of *Lafoensia pacari* and ellagic acid in a murine model of asthma. **Eur J Pharmacol** 508: 262 - 270.
- Rogério AP, Sá-Nunes A, Albuquerque DA, Anibal FF, Medeiros AI, Machado ER, Souza AO, Prado-Jr JC, Faccioli LH. 2003. *Lafoensia pacari* and IL-5 *Lafoensia pacari* extract inhibits IL-5 production in toxocariasis. **Parasite Immunol** 25: 393 - 400.
- Salgado HRN, Roncari AFF, Moreira RRD. 2005. Antidiarrhoeal effects of *Mikania glomerata* Spreng. (Asteraceae) leaf extract in mice. **Braz J Pharmacogn** 15: 205 - 208.
- Sampaio BL, Bara MTF, Ferri PH, Santos SC, Paula JR. 2011. Influence of environmental factors on the concentration of phenolic compounds in leaves of *Lafoensia pacari*. **Braz J Pharmacogn** 21: 1127 - 1137.
- Santos-Dantas V, Coelho-Dantas I, Pereira-Chaves T, Castro D, Silva H, Santos-Dantas GD. 2008. Análise das garrafadas indicadas pelos raizeiros na cidade de Campina Grande – PB. **Rev Biol Farm** 3: 33 - 50.

- Silva FS, Ramos MA, Hanazaki N, Albuquerque UP. 2011. Dynamics of traditional knowledge of medicinal plants in a rural community in the Brazilian semi-arid region. **Braz J Pharmacogn** 21: 382 - 391.
- Silva-Junior IF, Raimondi M, Zacchino S, Cechine-Filho V, Noldin VF, Rao VS, Lima JCS, Martins DTO. 2010. Evaluation of the antifungal activity and mode of action of *Lafoensia pacari* A. St.-Hil., Lythraceae, stem-bark extracts, fractions and ellagic acid. **Braz J Pharmacogn** 20: 422 - 428.
- Simões CM, Ribeiro-Do-Vale RM, Poli A, Nicolau M, Zanin M. 1989. The pharmacologic action of extracts of *Polygonum punctatum* Elliot (= *P. acre* HBK). **J Pharm Belg** 44: 275 - 284.
- Sivieri RRG, Guimarães PSS, Lima PDA, Neves GS, Graef CFF, Sivieri-Junior DO. 2015. **Atividade antioxidante do extrato da *Pseudobrickellia brasiliensis* Spreng.** Pôster na 67ª Reunião Anual da SBPC (Sociedade Brasileira para o Progresso da Ciência, São Carlos/SP, Brasil).
- Solon S, Lopes L, Teixeira-De-Sousa P, Schmeda-Hirschmann G. 2000. Free radical scavenging activity of *Lafoensia pacari*. **J Ethnopharmacol** 72: 173 - 178.
- Souza VC, Lorenzi H. 2012. **Botânica sistemática: guia ilustrado para identificação das famílias de fanerógamas nativas e exóticas no Brasil, baseado em APG III.** Instituto Plantarum, Nova Odessa/SP, Brasil.
- Souza, LKH, Oliveira CMA, Ferri PH, Santos SC, Oliveira-Júnior JG, Miranda ATB, Lião LM, Silva MRR. 2002. Antifungal properties of Brazilian Cerrado plants. **Braz J Microbiol** 33: 247 - 249.
- Tamashiro-Filho P, Sikiru OB, Tavares ADA, Lima JC, Marson-Ascêncio PG, Donizeti-Ascêncio S, Rios-Santos F, Martins DT. 2012. Evaluation of antiulcer activity and mechanism of action of methanol stem bark extract of *Lafoensia pacari* A. St.-Hil. (Lythraceae) in experimental animals. **J Ethnopharmacol** 144:497 - 505.
- Tan MJ, Ye JM, Turner N, Hohnen-Behrens C, Ke CQ, Tang CP, Chen T, Weiss HC, Gesing ER, Rowland A, James DE, Ye Y. 2008. Antidiabetic activities of triterpenoids isolated from bitter melon associated with activation of the AMPK pathway. **Chem Biol** 15: 263 - 273.
- Teixeira AS, Melo JIM. 2006. Plantas medicinais utilizadas no município de Jupi, Pernambuco, Brasil. **Iheringia Série Botânica** 61: 5 - 11.
- The Plant List. 2013. Version 1.1. Published on the Internet; <http://www.theplantlist.org>
- Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC, Erasmus BFN, Siqueira MF, Grainger A, Hannah L, Hughes L, Huntley B, van Jaarsveld AS, Midgley GF, Miles L, Ortega-Huerta MA, Peterson AT, Phillips OL, Williams SE. 2004. Extinction risk from climate change. **Nature** 427: 145 - 148.
- Tropicos. 2016. Tropicos.org, Missouri Botanical Garden. <http://www.tropicos.org>
- Valle TL. 2002. **Coleta de germoplasma de plantas cultivadas.** In: Amoroso MCM, Ming LC, Silva SP (eds.): Métodos de coleta e análise de dados em etnobiologia, etnoecologia e disciplinas correlatas. Ed. Unesp, Rio Claro/SP, Brasil.
- Valverde AL, Cardoso GL, Pereira NA, Silva AJ, Kuster RM. 2001. Analgesic and antiinflammatory activities of vernonioside B2 from *Vernonia condensata*. **Phytother Res** 15: 263 - 264.
- Vásquez SPF, Mendonça MSM, Noda SN. 2014. Etnobotânica de plantas medicinais em comunidades ribeirinhas do Município de Manacapuru, Amazonas, Brasil. **Acta Amaz** 44: 457 - 472.
- Vieira GT, Humberto JL, De Oliveira TT, Nagem TJ, Cecon PR, Braga FC. 2006. **Estudo do extrato butanólico de *Equisetum giganteum* L. na síndrome diabetes.** Anais da Reunião Anual da Sociedade Brasileira de Química.
- Vieira-Jr. HT, Moraes JM, Schobbenhaus C. 2011. **Geoparque Chapada dos Guimarães-MT: proposta.** Ed. CPRM, MT, Brasil.
- Zeni ALB, Bosio F. 2011. O uso de plantas medicinais em uma comunidade rural de Mata Atlântica - Nova Rússia, SC. **Neotrop Biol Conserv** 6: 55 - 63.